

# Surface Tension in Ising Systems with Kac Potentials

G. Alberti,<sup>1</sup> G. Bellettini,<sup>1</sup> M. Cassandro,<sup>2</sup> E. Presutti<sup>3</sup>

*Received January 6, 1995*

---

We consider an Ising spin system with Kac potentials in a torus of  $\mathbb{Z}^d$ ,  $d \geq 2$ , and fix the temperature below its Lebowitz-Penrose critical value. We prove that when the Kac scaling parameter  $\gamma$  vanishes, the log of the probability of an interface becomes proportional to its area and the surface tension, related to the proportionality constant, converges to the van der Waals surface tension. The results are based on the analysis of the rate functionals for Gibbsian large deviations and on the proof that they  $\Gamma$ -converge to the perimeter functional of geometric measure theory (which extends the notion of area). Our considerations include non smooth interfaces proving that the Gibbsian probability of an interface depends only on its area and not on its regularity.

---

**KEY WORDS:** Interfaces; Kac potentials, surface tension,  $\Gamma$ -convergence.

## 1. INTRODUCTION

The thermodynamic free energy excess of a simple fluid when two pure phases coexist is

$$F = \int_{\Sigma} d\lambda(r) s_{\beta}(n(r)) \quad (1.1)$$

where  $\Sigma$  is the interface that separates the two phases,  $d\lambda(r)$  is the surface area element, and  $n(r)$  is the unit normal to  $\Sigma$  at  $r$ ; finally,  $s_{\beta}(n)$  is the surface tension at the inverse temperature  $\beta$  of a flat interface perpendicular

---

<sup>1</sup>Istituto di Matematiche Applicate "U. Dini", 56126 Pisa, Italy.

<sup>2</sup>Dipartimento di Fisica, Università di Roma La Sapienza, 00185 Rome, Italy.

<sup>3</sup>Dipartimento di Matematica, Università di Roma Tor Vergata, 00133 Rome, Italy.

to  $n$ . The purpose of this paper is to derive (1.1) in the context of Ising spin systems with Kac potentials. In particular we compute the surface tension, proving the validity of the expression proposed by the van der Waals theory.

In this introduction we stress the more physical aspects, leaving the mathematical details to the next sections. The surface tension is usually defined in Ising systems by a formula like

$$s_{\beta,\Lambda} := \frac{1}{\beta|\partial\Lambda|} \log \frac{Z_{\beta,\Lambda}^{++}}{Z_{\beta,\Lambda}^{+-}} \quad (1.2)$$

where  $\Lambda$  is (for instance) a cube in  $\mathbb{R}^d$  with unit vector  $n$  normal to its top face,  $|\partial\Lambda|$  being the corresponding area;  $Z_{\beta,\Lambda}^{++}$  and  $Z_{\beta,\Lambda}^{+-}$  are then the partition functions with  $++$  and  $+-$ , respectively; boundary conditions on the top and the bottom of  $\Lambda$  and periodic conditions on the other sides.

The relation with (1.1) comes from the assumption that the main contribution to the free energy difference when changing  $+$  into  $-$  in the bottom is due to the appearance of a flat interface normal on the average to  $n$ . According to (1.1), this excess free energy should then be  $|\partial\Lambda|s_{\beta}(n)$ , which gives (1.2), but an exact equality can be achieved only in the thermodynamic limit where fluctuations are depressed. The existence of this thermodynamic limit has been proved for several classes of models; see, for instance, Bricmont *et al.* [2]. A definition of the surface tension in terms of (1.2) is thus based on a preliminary assumption of the validity of (1.1), which conceptually should be derived first.

Let us now turn to the probabilistic aspects of the issue, which lead us to large deviations, as can be most clearly seen in a formulation where (1.1) appears again in a somewhat oblique fashion. Call  $m_{\Lambda}$  the empirical average (i.e., the total magnetization density) of the spins in the cube  $\Lambda$ , where the Gibbs measure with periodic boundary conditions is defined. In our context the Gibbs distribution of  $m_{\Lambda}$  when  $\Lambda$  invades the whole space becomes supported by two values that we call  $\pm m_{\beta}$  (in the case we consider there is a spin flip symmetry). Thus  $\pm m_{\beta}$  are the pure phase equilibrium magnetizations. The analysis of the distribution of  $m_{\Lambda}$  away from  $\pm m_{\beta}$  is a large-deviation problem. Remarkably, in Ising systems in  $d = 2$  dimensions and nearest neighbor ferromagnetic interactions there is a complete answer for all temperatures below the critical one [15, 27]. The probability that  $m_{\Lambda}$  is “close” to a value  $m$  in  $(-m_{\beta}, m_{\beta})$  is found to vanish as  $\exp(-c|\partial\Lambda|)$  when  $|\Lambda| \rightarrow +\infty$ . The rate  $c$  comes from the solution (Wulff Construction) of a variational problem involving (1.1) whose validity is an indirect consequence of the proof.

A direct proof of (1.1) is the true goal of this paper. First of all we need a well-defined setting of the problem which involves an interpretation of (1.1) as a functional in an appropriate function space. To this end it is better to regard  $F$  as a function of the magnetization profile that has  $\Sigma$  as its interface. We thus consider functions  $u(r)$ ,  $r \in \mathcal{T}$ , with only two values  $\pm m_\beta$ . Then  $\Sigma$  is defined as the boundary of the set  $\{r \in \mathcal{T} : u(r) = +m_\beta\}$ . The minimal requirement on  $u$  for (1.1) to hold is that this should be finite. A general notion of area has been developed in “geometric measure theory” (see, for instance, [6], [17], [18], [20], [30]), where it is defined as a functional  $P(u)$  with  $u$  (in our context) in  $BV(\mathcal{T}; \{\pm m_\beta\})$ . Then  $P(u)$  generalizes the classical notion of area of the interface of  $u$  and it is finite on each element of  $BV(\mathcal{T}; \{\pm m_\beta\})$ . On such a space the formula (1.1) is well defined with  $d\lambda(r)$  the Hausdorff measure on the “reduced boundary” of  $\Sigma$  [17], provided  $s_\beta(n)$  is measurable on the unit sphere of  $\mathbb{R}^d$ .

While smooth bounded surfaces are included, in this class there are also highly irregular surfaces with finite area. The choice of the domain where  $F$  is defined therefore has important implications in the derivation of (1.1), namely whether it is the area the only factor that determines the probability of an interface or there are other features such as the regularity of the surface which play an effective role. We will see that for Ising systems with Kac potentials the area alone determines the probability of an interface.

The next question concerns the quantity which should play the statistical mechanical role of the functional (1.1). Since  $u(r)$  is a “macroscopic density magnetization profile”, the relevant quantity is the Gibbsian probability of “recognizing such a profile” out of the actual spin configurations. The region  $\Lambda$  where the Ising spins are defined should then be scaled down to a fixed region to be identified with the above unit torus  $\mathcal{T}$  where spin configurations will be represented in terms of piecewise constant functions with values  $\pm 1$ ; precise definitions are given in the next section. The Gibbs measures can be regarded as probabilities on  $L^\infty(\mathcal{T}; [-1, 1])$  and we are interested in computing the probability of sets  $A_\zeta(u)$  which are neighborhoods of  $u \in BV(\mathcal{T}; \{\pm m_\beta\})$  (in the  $L^1(\mathcal{T}; [-1, 1])$ -topology, as justified in the next section) with “width” determined by  $\zeta$ , a positive parameter. Calling  $M_{\beta, \varepsilon, \zeta}(u)$  such a probability, where  $\beta$  is the inverse temperature (of the Gibbs measure) and  $\varepsilon$  the scaling parameter which gives the ratio between microscopic and macroscopic units, the quantity that approximates  $F$  in (1.1) should then be

$$\frac{\varepsilon^{d-1}}{\beta} \log M_{\beta, \varepsilon, \zeta}(u) \tag{1.3}$$

in the limit where first  $\varepsilon \rightarrow 0^+$  (thermodynamic limit) and then  $\zeta \rightarrow 0^+$ , i.e.,

the limit of high accuracy in the recognition of  $u$ . The quantity  $\varepsilon^{d-1}$  scales as a surface and plays the role of  $|\partial\Lambda|^{-1}$  in the previous considerations.

The quantity (1.3) has to be compared with the previous large-deviation probabilities, and in fact  $F$  in (1.1) should be regarded as the rate function of large deviations associated with (1.3). It is, however, convenient at this point to particularize the discussion to the case of the Ising ferromagnetic Kac potentials [22] that we actually study in this paper. Here we have another parameter  $\gamma > 0$  that determines the range of the interaction. One is interested in the limit as  $\gamma \rightarrow 0^+$ , where the range of the interaction diverges as  $\gamma^{-1}$ , as recalled in the next section. We will study the simultaneous limit where  $\varepsilon \rightarrow 0^+$  (thermodynamic limit) and  $\gamma \rightarrow 0^+$  (scaling limit) and for technical reasons we will ask for a strict relation between the two. One of the features of the Kac potentials is that large-deviation rate functional as  $\gamma \rightarrow 0^+$  has an explicit form in terms of the nonlocal van der Waals functional that we called  $F_\varepsilon(u)$ ,  $u \in L^\infty(\mathcal{I}, [-1, 1])$  ( $\beta$  is not made explicit here). According to this statement we can replace (in the limit when  $\varepsilon$  and  $\gamma$  vanish) (1.3) by

$$\inf_{v \in A_\zeta(u)} F_\varepsilon(v) \tag{1.4}$$

The validity of (1.1) is then in this setup just the statement that  $F(u)$  in (1.1) is the limit of (1.4) as  $\varepsilon \rightarrow 0^+$  and then  $\zeta \rightarrow 0^+$ , which on the other hand is exactly the setting of De Giorgi's notion of  $\Gamma$ -convergence (of  $F_\varepsilon$  to  $F$ ) [4, 7].

We actually prove (see Theorem 2.5) that  $F_\varepsilon$   $\Gamma$ -converge to the perimeter functional  $P(u)$ , up to a multiplicative factor  $s_\beta$ , which is then the isotropic surface tension of the system. In agreement with the van der Waals theory [see (2.22)],  $s_\beta$  can be expressed as the free energy of the instanton solution of a nonlocal Euler-Lagrange equation for the functionals  $F_\varepsilon$ .

Let us briefly describe the content of the paper. In Section 2 we state the main definitions and results; we also outline the proofs, whose details are reported in the next two sections and in the appendices. In Section 3 we show how the probability estimates involving the Gibbs measures are expressed in terms of the functional  $F_\varepsilon$  and the validity of (1.1) related to a problem of  $\Gamma$ -convergence, which is solved in Section 4; more technical problems are left to the appendices. As the relation with the  $\Gamma$ -convergence of  $F_\varepsilon$  may have interest in its own right, we have isolated the whole argument: the problem of  $\Gamma$ -convergence is formulated at the end of Section 2 and studied in Section 4, which can be read independently of Section 3.

In a forthcoming paper [1], the above results are extended to include an analysis of the Wulff and other constrained variational problems. It is also

proved that the interfaces with infinite area have superexponentially small probability.

## 2. MAIN RESULT

We start by recalling the notion of Ising systems with Kac potentials. Maybe our notation will not be the most usual ones, see the Remarks after Definition 2.1d below, but this is going to be the most convenient setup for our analysis. We split the main definition into several ones.

**Definition 2.1a.** *Partitions of  $\mathbb{R}^d$ .* For any  $k \in \mathbb{Z}$ ,  $\mathcal{Q}^{(k)}$  denotes the partition of  $\mathbb{R}^d$  into the  $d$ -dimensional cubes

$$\{r = (r_1, \dots, r_d) \in \mathbb{R}^d : 2^{-k}x_i \leq r_i < 2^{-k}(x_i + 1); x_i \in \mathbb{Z}, i = 1, \dots, d\} \quad (2.1)$$

The atoms of  $\mathcal{Q}^{(k)}$  are denoted by  $C^{(k)}$ .  $C^{(k)}(r)$  is the unique atom of  $\mathcal{Q}^{(k)}$  that contains the point  $r$ . A function  $f \in L^\infty(\mathbb{R}^d)$  is  $\mathcal{Q}^{(k)}$ -measurable if it is constant on the atoms of  $\mathcal{Q}^{(k)}$  and a set  $A \subset \mathbb{R}^d$  is  $\mathcal{Q}^{(k)}$ -measurable if its characteristic function  $\mathbf{1}_A$  is  $\mathcal{Q}^{(k)}$ -measurable.

**Definition 2.1b.** *Spin configurations.* We denote by  $\gamma$  a parameter that takes values in  $\{2^{-k}, k \in \mathbb{N}\}$ . Let  $\gamma = 2^{-k_\gamma}$ ,  $k_\gamma \in \mathbb{N}$ , and  $\varepsilon^{-1} \in \mathbb{N}$ ; we say that  $\sigma_\gamma$  is a spin configuration with mesh  $\gamma$  and period  $\varepsilon^{-1}$  if  $\sigma_\gamma \in L^\infty(\mathbb{R}^d; \{\pm 1\})$ ,  $\sigma_\gamma$  is  $\mathcal{Q}^{(k_\gamma)}$ -measurable and if

$$\sigma_\gamma(r) = \sigma_\gamma(r') \text{ whenever } r_i - r'_i = \varepsilon^{-1}x_i, \text{ where } x_i \in \mathbb{Z} \text{ for } i = 1, \dots, d$$

Denoting by  $\mathcal{T}_\varepsilon$  the torus in  $\mathbb{R}^d$  of period  $\varepsilon^{-1}$ , a spin configuration on  $\mathcal{T}_\varepsilon$  will be identified with its  $\varepsilon^{-1}$ -periodic extension to  $\mathbb{R}^d$ .

The values of  $\sigma_\gamma$  in each atom of  $\mathcal{Q}^{(k)}$  are the spins of the configuration  $\sigma_\gamma$ .

**Definition 2.1c.** *Energy.* The interaction strength  $J(|r|)$  is a non-negative  $\mathcal{C}^\infty$  function of  $r \in \mathbb{R}^d$ , supported in the unit ball, with  $\sup\{s > 0 : J(s) > 0\} = 1$  and such that

$$\int_{\mathbb{R}^d} dr J(|r|) = 1 \quad (2.2)$$

Let  $\Lambda$  be a bounded measurable region in  $\mathbb{R}^d$  and  $m \in L^\infty(\Lambda; [-1, 1])$ . The energy of  $m$  in  $\Lambda$  is defined as

$$H(m; \Lambda) := -\frac{1}{2} \int_{\Lambda} dr \int_{\Lambda} dr' J(|r - r'|) m(r) m(r') \quad (2.3)$$

If  $\Lambda$  is a torus then  $|r - r'|$  in (2.3) is the distance between  $r$  and  $r'$  in the torus.

The ferromagnetic condition  $J \geq 0$  will be essential in most of the proofs.

**Definition 2.1d.** *Gibbs measure.* The Gibbs measure on the torus  $\mathcal{T}_\varepsilon$ , with Kac potential  $J(|r|)$ , scaling parameter  $\gamma$  and inverse temperature  $\beta$  is the probability  $\mu_{\beta,\gamma,\varepsilon}$  on the space of spin configurations on  $\mathcal{T}_\varepsilon$  with mesh  $\gamma$  defined as

$$\mu_{\beta,\gamma,\varepsilon}(\sigma_\gamma) := \frac{1}{Z_{\beta,\gamma,\varepsilon}} \exp[-\beta\gamma^{-d}H(\sigma_\gamma; \mathcal{T}_\varepsilon)] \quad (2.4)$$

where

$$Z_{\beta,\gamma,\varepsilon} := \sum_{\sigma_\gamma} \exp[-\beta\gamma^{-d}H(\sigma_\gamma; \mathcal{T}_\varepsilon)]. \quad (2.5)$$

is the partition function.

**Remarks.** Calling

$$S(x) := \sigma_\gamma(\gamma x), \quad x \in \mathbb{Z}^d$$

and  $T_\varepsilon := \{x \in \mathbb{Z}^d : \gamma x \in \mathcal{T}_\varepsilon\}$ , we have

$$\gamma^{-d}H(\sigma_\gamma; \mathcal{T}_\varepsilon) = -\frac{1}{2} \sum_{\substack{x,y \in T_\varepsilon \\ x \neq y}} J_\gamma(x,y) S(x)S(y) + c_\varepsilon \quad (2.6)$$

where  $c_\varepsilon$  (which takes into account the sum over  $x = y$ ) is independent of  $\sigma_\gamma$ , and, recalling  $\gamma = 2^{-k_\gamma}$ ,

$$J_\gamma(x,y) := \gamma^{-d} \int_{C^{(k_\gamma)}(\gamma x)} dr \int_{C^{(k_\gamma)}(\gamma y)} dr' J(|r - r'|) \quad (2.7)$$

Observe that the coefficient  $c_\varepsilon$  drops from the expression for the Gibbs measure and it is therefore irrelevant. Then, using the variables  $S(x)$ , the energy and the Gibbs formula take the usual form, in particular neglecting the variation of  $J$  in the integral in (2.7) we get

$$J_\gamma(x,y) \approx \gamma^d J(\gamma|x - y|) \quad (2.8)$$

which has the typical scaling properties of the original Kac potential. To simplify the notation we have directly defined the model with the interaction

(2.7), but the results in this paper hold as well when the energy is given by (2.6) with (2.8) holding as an equality.

The system in Definition 2.1 is thus included in the class introduced by Kac to model the van der Waals theory of phase transition, which is in fact derived by taking the limit  $\gamma \rightarrow 0^+$  [22]. The physically correct procedure would be to let first  $\varepsilon^{-1} \rightarrow +\infty$  and then  $\gamma \rightarrow 0^+$ , but we will instead study the much simpler problem where  $\varepsilon^{-1}$  diverges “not too fast” as  $\gamma \rightarrow 0^+$ , see (2.10) below. As we shall see even, in this regime there are interesting phenomena.

**Definition 2.1e.** *Choice of parameters.* In the sequel we fix  $\beta > 1$  and, setting

$$0 < \alpha < \frac{1}{d+1} \quad (2.9)$$

we choose

$$\varepsilon^{-1} := [\gamma^{-\alpha}] \quad (2.10)$$

where  $[a]$  denotes the integer part of  $a$ .

The system with  $\beta > 1$  has a phase transition when  $\gamma \rightarrow 0^+$ , that is in the Lebowitz-Penrose limit [23]. There are two equilibrium magnetizations,  $\pm m_\beta$ , where

$$m_\beta = \tanh\{\beta m_\beta\}, \quad m_\beta > 0 \quad (2.11)$$

(which has a solution  $0 < m_\beta < 1$  if and only if  $\beta > 1$ ). The equilibrium magnetizations are defined in [23] in terms of the partition function. We will extend the result by showing that  $\pm m_\beta$  are also the magnetization densities of the typical Gibbs spin configurations. We will in fact prove in Theorem 2.3 that the probability of the configurations which have either magnetization  $m_\beta$  or  $-m_\beta$  converges in the limit  $\gamma \rightarrow 0^+$  to 1. We will then investigate the residual configurations, in particular those which have an interface. Since by Theorem 2.3 they have vanishing probability, this will be a problem of large deviations, but, as we shall see [Eq. (2.16)], with an “anomalous normalization”.

For any  $r \in \mathbb{R}^d$  and  $R > 0$  we set  $B_R(r) = \{r' \in \mathbb{R}^d : |r - r'| \leq R\}$  and use the shorthand notation

$$\int_{B_R(r)} dr' f(r') = \frac{1}{|B_R(r)|} \int_{B_R(r)} dr' f(r') \quad (2.12)$$

**Definition 2.2.** Given  $-1 \leq m \leq 1$ ,  $R > 0$  and  $\zeta > 0$ , we say that a spin configuration  $\sigma_\gamma$  on  $\mathcal{T}_\varepsilon$  has magnetization constantly equal to  $m$  with

accuracy  $(R, \zeta)$  if

$$\varepsilon^d \int_{\mathcal{T}_\varepsilon} dr \left| m - \int_{B_R(r)} dr' \sigma_\gamma(r') \right| < \zeta \quad (2.13)$$

and write  $\sigma_\gamma \in \mathcal{P}_{R, \zeta, \gamma}^\pm$  if  $\sigma_\gamma$  satisfies (2.13) with  $m = \pm m_\beta$ .

It is clearly necessary to define the magnetization of a spin configuration via an averaging procedure because the spins have only values  $\pm 1$ . There is, however, some degree of arbitrariness about the size of the averaging region, we have chosen regions with finite volumes (in interaction range units). Observe that (2.13) does not imply that the averages are uniformly close to  $m$ , but that this only happens in a large fraction of the whole volume. Such a weaker condition is more likely to extend to systems where the condition that  $\alpha$  in (2.9) is small is either relaxed or dropped.

In the next Section we will prove the following result.

**Theorem 2.3** Let  $\alpha, \varepsilon$  be as in (2.9), (2.10) respectively, and let  $R > 0$ . Then there is  $\zeta_\gamma \rightarrow 0^+$  as  $\gamma \rightarrow 0^+$  so that

$$\lim_{\gamma \rightarrow 0^+} \mu_{\beta, \gamma, \varepsilon}(\mathcal{P}_{R, \zeta_\gamma, \gamma}^- \cup \mathcal{P}_{R, \zeta_\gamma, \gamma}^+) = 1. \quad (2.14)$$

In the complement of  $\mathcal{P}_{R, \zeta_\gamma, \gamma}^- \cup \mathcal{P}_{R, \zeta_\gamma, \gamma}^+$  there are configurations that describe coexistence of phases and interfaces, to see this we need to generalize Definition 2.2, replacing the number  $m$  by a function  $m(\cdot)$ . Denoting by  $\mathcal{T}$  the unit torus in  $\mathbb{R}^d$  and by  $L^\infty(\mathcal{T}; [-1, 1])$  the space of integrable functions on  $\mathcal{T}$  with values in  $[-1, 1]$ , we set:

**Definition 2.4.** Let  $m \in L^\infty(\mathcal{T}; [-1, 1])$ , and  $R, \zeta \in \mathbb{R}^+$ . We say that a spin configuration  $\sigma_\gamma$  on  $\mathcal{T}_\varepsilon$  has magnetization  $m$  with accuracy  $(R, \zeta)$  if  $\sigma_\gamma \in \mathcal{P}_{R, \zeta, \gamma}(m)$  where

$$\mathcal{P}_{R, \zeta, \gamma}(m) = \left\{ m^* \in L^\infty(\mathcal{T}_\varepsilon; [-1, 1]) : \varepsilon^d \int_{\mathcal{T}_\varepsilon} dr \left| \int_{B_R(r)} dr' (m(\varepsilon r') - m^*(r')) \right| < \zeta \right\} \quad (2.15)$$

Let  $m \in L^\infty(\mathcal{T}; \{\pm m_\beta\})$ , we then call the regions  $\{r \in \mathcal{T} : m(r) = \pm m_\beta\}$  as the  $\pm$  phases of  $m$  and define the interface of  $m$  as the boundary of the plus-phase. On physical grounds the cost in free energy to create an interface is proportional to its area, the proportionality constant being the surface tension. Usually the surface tension is defined by imposing plus and



minus conditions at the top and the bottom of a rectangular region, see for instance [2], so that the interface is (in the average) flat and parallel to the bases of the rectangle. We will prove that the free energy is still proportional to the area times the surface tension even when the surface is not regular. Let  $P(m)$  be the perimeter functional on  $BV(\mathcal{I}; \{\pm m_\beta\})$  (the functions of bounded variation on  $\mathcal{I}$  with values  $\pm m_\beta$ ) that defines the area of the plus-phase of  $m$ , see Appendix D. Then:

**Theorem 2.5.** There is  $s_\beta > 0$ , given in (2.22) below, so that the following holds. For all  $u \in BV(\mathcal{I}; \{\pm m_\beta\})$  and all  $R > 0$  there is  $\zeta_\gamma \rightarrow 0^+$  as  $\gamma \rightarrow 0^+$  so that

$$\lim_{\gamma \rightarrow 0^+} -\gamma^d \varepsilon^{d-1} \log[\mu_{\beta, \gamma, \varepsilon}(\mathcal{P}_{R, \zeta_\gamma, \gamma}(u))] = \beta s_\beta P(u) \quad (2.16)$$

**Remarks.** The number of spins in a spin configuration is proportional to  $(\gamma\varepsilon)^{-d}$ . Then, “in normal conditions”, the large deviation normalization factor is  $(\gamma\varepsilon)^d$ . In this sense the normalization in (2.16) is anomalous, the anomaly is due to the presence of a phase transition. Since there are two equilibrium magnetization values,  $\pm m_\beta$ , it is possible to have a non constant profile with equilibrium magnetization at all points: such profiles only cost a surface price. To get the prefactor  $\gamma^d \varepsilon^{d-1}$  recall that the surfaces scale like  $\varepsilon^{1-d}$ , as  $\mathcal{I}_\varepsilon = \varepsilon^{-1} \mathcal{I}$ . As shown in the sequel, the thickness of the interface (in  $\mathcal{I}_\varepsilon$ ) is of the order of unity, thus the volume of the region around the interface scales also as  $\varepsilon^{1-d}$ . The number of spins in a region is proportional to the volume of the region times  $\gamma^{-d}$ , we thus obtain  $\gamma^{-d} \varepsilon^{1-d}$  which is the inverse of the prefactor in (2.16).

To give an expression to  $s_\beta$  we need several intermediate definitions. We start by recalling a result proved in [14]:

**Theorem 2.6.** There is a unique function  $\bar{m} : \mathbb{R} \rightarrow [-1, 1]$  such that  $\bar{m}(0) = 0$ ,

$$\liminf_{s \rightarrow +\infty} \bar{m}(s) > 0, \quad \limsup_{s \rightarrow -\infty} \bar{m}(s) < 0$$

and such that for all  $s \in \mathbb{R}$

$$\bar{m}(s) = \tanh\{\beta \bar{J} * \bar{m}(s)\} \quad (2.17)$$

where  $*$  denotes convolution and for every  $s \in \mathbb{R}$

$$\bar{J}(s) := \int_{\mathbb{R}^{d-1}} dr J((s^2 + r^2)^{1/2}) \quad (2.18)$$

Moreover  $\bar{m} \in \mathcal{C}^\infty(\mathbb{R})$ , it is antisymmetric, strictly increasing and with asymptotic values at  $\pm\infty$  equal to  $\pm m_\beta$ , to which it converges exponentially fast.

As we shall see  $\bar{m}$ , called the instanton, describes the magnetization pattern at the interface. We next define the excess free energy functional  $\mathcal{F}$ :

**Definition 2.7.** Given a measurable set  $\Lambda \subset \mathbb{R}^d$  and  $m \in L^\infty(\Lambda; [-1, 1])$  we define the map  $\mathcal{F}(m; \Lambda)$  with values in  $[0, +\infty]$  as

$$\begin{aligned} \mathcal{F}(m; \Lambda) &:= \int_{\Lambda} dr [f(m(r)) - f(m_\beta)] + \\ &+ \frac{1}{4} \iint_{\Lambda \times \Lambda} dr dr' J(|r - r'|) [m(r) - m(r')]^2 \end{aligned} \quad (2.19)$$

where, if  $s \in \mathbb{R}$ ,  $f(s)$  and  $i(s)$  are the free energy and the entropy density, namely

$$f(s) := -\frac{1}{2}s^2 - \beta^{-1}i(s) \quad (2.20)$$

$$i(s) := -\frac{1+s}{2} \log\left(\frac{1+s}{2}\right) - \frac{1-s}{2} \log\left(\frac{1-s}{2}\right) \quad (2.21)$$

If  $\Lambda$  is a torus,  $|r - r'|$  is the distance between  $r$  and  $r'$  in the torus. We further set  $\mathcal{F}(m) := \mathcal{F}(m; \mathbb{R}^d)$  and  $\mathcal{F}_\varepsilon(m) := \mathcal{F}(m; \mathcal{I}_\varepsilon)$ .

We call  $\bar{\mathcal{F}}(m)$  the  $d = 1$  version of  $\mathcal{F}(m)$  with  $J$  replaced by  $\bar{J}$ , see (2.18). Together with Theorem 2.5 we shall prove that

$$s_\beta = \bar{\mathcal{F}}(\bar{m}) \quad (2.22)$$

The proof of Theorem 2.5 starts from a large deviation estimate for the Gibbs measure. Because of the assumption on the size of the region we are essentially reduced to the case considered by Eisele and Ellis, [16], and the large deviation rate function is the functional  $\mathcal{F}_\varepsilon$ .

Given  $m \in L^\infty(\mathcal{I}_\varepsilon; [-1, 1])$  we set

$$u(r) := m(\varepsilon^{-1}r), \quad r \in \mathcal{I}$$

and define the functional

$$F_\varepsilon(u) := \varepsilon^{d-1} \mathcal{F}_\varepsilon(m) \quad (2.23)$$

on  $L^\infty(\mathcal{I}; [-1, 1])$ , thus

$$F_\varepsilon(u) = \varepsilon^{-1} \int_{\mathcal{I}} dr [f(u(r)) - f(m_\beta)] + \frac{\varepsilon}{4} \iint_{\mathcal{I} \times \mathcal{I}} dr dr' J_\varepsilon(|r - r'|) \left[ \frac{u(r) - u(r')}{\varepsilon} \right]^2, \quad (2.24)$$

where

$$J_\varepsilon(|r|) := \varepsilon^{-d} J(\varepsilon^{-1}|r|). \quad (2.25)$$

In the next Section we will prove that Theorem 2.5 follows by proving that  $\{F_\varepsilon\}$   $\Gamma$ -converges to  $F(u) := s_\beta P(u)$  as  $\varepsilon \rightarrow 0^+$ . This means that given any  $u \in BV(\mathcal{I}; \{\pm m_\beta\})$  the following holds.

1. For any family  $\{u_\varepsilon\} \subseteq L^\infty(\mathcal{I}; [-1, 1])$  that converges to  $u$  in  $L^1(\mathcal{I}; [-1, 1])$  as  $\varepsilon \rightarrow 0^+$  we have

$$\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) \geq F(u); \quad (2.26)$$

2. there exists a sequence  $\{u_\varepsilon\} \subseteq L^\infty(\mathcal{I}; [-m_\beta, m_\beta])$  that converges to  $u$  in  $L^1(\mathcal{I}; [-1, 1])$  as  $\varepsilon \rightarrow 0^+$  and such that

$$\lim_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) = F(u) \quad (2.27)$$

Since  $J_\varepsilon$  in (2.25) is an approximated delta function it might look reasonable to replace the second integral in (2.24) by

$$D \frac{\varepsilon}{4} \int_{\mathcal{I}} dr |\nabla u|^2, \quad \text{where } D := \int_{\mathbb{R}^d} dr J(|r|) r^2. \quad (2.28)$$

We then obtain the classical example of functionals that  $\Gamma$ -converge to  $P(u)$  (modulo the constant factor  $\sqrt{D}/2$ ), as conjectured by De Giorgi in 1975 [7] and proved by Modica and Mortola, [26]. However the constant is not the surface tension of our model!

In Section 4 we will prove that  $\{F_\varepsilon\}$   $\Gamma$ -converges to  $s_\beta P(u)$ .

### 3. REDUCTION TO A VARIATIONAL PRINCIPLE

In the first part of this Section we prove a relation between the Gibbs probability  $\mu_{\beta, \gamma, \varepsilon}$  and the functional  $\mathcal{F}_\varepsilon(m)$  by showing that, for small  $\gamma$ ,  $\mu_{\beta, \gamma, \varepsilon}$  is well approximated by

$$\mu_{\beta, \gamma, \varepsilon}(\cdot) \approx \exp[-\beta \gamma^{-d} \mathcal{F}_\varepsilon(\cdot)] \quad (3.1)$$

The sense in which (3.1) holds is specified in Lemma 3.2 below. We can already say however that (3.1) will not refer to single spin configurations for which it is not valid, but rather to a coarse grained version of the configuration itself that we will define in the sequel.

In the second part of the Section we will see, with the help of (3.1), that the problem of computing a probability may be reduced to finding minima of the functional  $\mathcal{F}_\varepsilon$ . In this way we will relate the proof of Theorem 2.5 to the  $\Gamma$ -convergence problem stated at the end of Section 2.

We begin with a few definitions aimed at introducing the basic notion of coarse grained configurations.

### 3.1. The Partition $\mathcal{Q}_\gamma$

Recalling the definition of  $\alpha$  in (2.9) we choose  $\delta > 0$  so that

$$d\alpha < \delta < 1 - \alpha \quad (3.2)$$

and call  $\mathcal{Q}_\gamma$  the partition  $\mathcal{Q}^{(k)}$  of Definition 2.1a with  $k = n_\gamma$  and

$$n_\gamma := \left\lceil \frac{\delta \log(\gamma^{-1})}{\log 2} \right\rceil, \quad \text{so that } \gamma^\delta \leq 2^{-n_\gamma} < 2\gamma^\delta \quad (3.3)$$

### 3.2. The Coarse Grained Configurations

Recalling that  $C^{(k)}(r)$  is the atom of  $\mathcal{Q}^{(k)}$  that contains  $r$ , we let  $\pi^{(k)}$  be the map from  $L^\infty(\mathbb{R}^d)$  into itself defined by

$$\pi^{(k)} f(r) := \frac{1}{|C^{(k)}|} \int_{C^{(k)}(r)} dr' f(r') \quad (3.4)$$

We then set  $\pi_\gamma := \pi^{(n_\gamma)}$ ,  $s^{(k)} := \pi^{(k)} \sigma_\gamma$  and  $s_\gamma := \pi_\gamma \sigma_\gamma$ , where  $\sigma_\gamma$  is a spin configuration, Definition 2.1b;  $s_\gamma$  will be called a coarse grained (spin) configuration

### 3.3. Weights and Energies of the Coarse-Grained (Spin) Configurations

The weight  $W(s_\gamma)$  of the coarse grained configuration  $s_\gamma$  is the number of spin configurations  $\sigma_\gamma$  such that  $\pi_\gamma \sigma_\gamma = s_\gamma$ .

We denote by  $H_\varepsilon(\cdot) := H(\cdot; \mathcal{T}_\varepsilon)$ , see (2.3). The energy of a coarse grained configuration  $s_\gamma$  is then denoted by  $H_\varepsilon(s_\gamma) = H(s_\gamma; \mathcal{T}_\varepsilon)$ .

We finally write, by an abuse of notation,

$$\mu_{\beta, \gamma, \varepsilon}(s_\gamma) = \mu_{\beta, \gamma, \varepsilon}(\{\sigma_\gamma : \pi_\gamma \sigma_\gamma = s_\gamma\}) \quad (3.5)$$

namely  $\mu_{\beta, \gamma, \varepsilon}(s_\gamma)$  is the probability of all the spin configurations  $\sigma_\gamma$  whose coarse grained image is  $s_\gamma$ .

In the next Lemma we relate the energy and the weight of the coarse grained configurations respectively to the original energy of the spin configurations and to the entropy functional  $I$ . The latter is defined, for any given bounded measurable region  $\Lambda$  in  $\mathbb{R}^d$ , as the functional  $I(m; \Lambda)$  on  $L^\infty(\Lambda; [-1, 1])$ :

$$I(m; \Lambda) := \int_\Lambda dr i(m(r)) \quad (3.6)$$

with  $i(m)$  as in (2.21). We set  $I_\varepsilon(m) := I(m; \mathcal{T}_\varepsilon)$ .

**Lemma 3.1.** There are positive constants  $c_1, c_2, c_3$  and  $c_4$  such that for all spin configurations  $\sigma_\gamma$ , all  $r \in \mathbb{R}^d$  and all  $R > 0$

$$\left| \int_{B_R(r)} dr' \sigma_\gamma(r') - \int_{B_R(r)} dr' \pi_\gamma \sigma_\gamma(r') \right| \leq c_1 R^{-1} \gamma^\delta \quad (3.7)$$

Moreover

$$|H_\varepsilon(\sigma_\gamma) - H_\varepsilon(\pi_\gamma \sigma_\gamma)| \leq c_2 \gamma^\delta \varepsilon^{-d} \quad (3.8)$$

and for all coarse grained configurations  $s_\gamma$

$$|\log W(s_\gamma) - \gamma^{-d} I_\varepsilon(s_\gamma)| \leq c_3 (\gamma \varepsilon)^{-d} \gamma^{2(1-\delta)d} \log(\gamma^{-1}) \quad (3.9)$$

Finally, denoting by  $N_\gamma$  the total number of distinct coarse grained configurations  $s_\gamma$ :

$$\log N_\gamma \leq c_4 (\gamma \varepsilon)^{-d} \gamma^{(1-\delta)d} \log(\gamma^{-1}) \quad (3.10)$$

We prove the lemma in Appendix A.

The functionals  $I_\varepsilon$  and  $H_\varepsilon$  are related to  $\mathcal{F}_\varepsilon$  in a simple way: by expanding the square in the last term of (2.19), recalling the definition (2.20) of  $f(m)$  and using (2.2), we get

$$\begin{aligned} \mathcal{F}_\varepsilon(m) &= [H_\varepsilon(m) - \beta^{-1} I_\varepsilon(m)] - [H_\varepsilon(\hat{m}_\beta) - \beta^{-1} I_\varepsilon(\hat{m}_\beta)] \\ &\quad \text{where } \hat{m}_\beta(r) \equiv m_\beta, r \in \mathcal{T}_\varepsilon \end{aligned} \quad (3.11)$$

Then, as a corollary of Lemma 3.1, we have the following version of (3.1).

**Lemma 3.2.** There is a constant  $C_1 > 0$  so that for any coarse grained spin configuration  $s_\gamma$

$$|\log \mu_{\beta,\gamma,\varepsilon}(s_\gamma) + \beta\gamma^{-d}\mathcal{F}_\varepsilon(s_\gamma)| \leq C_1(\gamma\varepsilon)^{-d}[\gamma^{(1-\delta)d}\log(\gamma^{-1}) + \gamma^\delta] \quad (3.12)$$

*Proof.* By (3.5) and (2.4)

$$\mu_{\beta,\gamma,\varepsilon}(s_\gamma) = \sum_{\pi_\gamma \sigma_\gamma = s_\gamma} \frac{1}{Z_{\beta,\gamma,\varepsilon}} \exp[-\beta\gamma^{-d}H_\varepsilon(\sigma_\gamma)]$$

where  $Z_{\beta,\gamma,\varepsilon}$  denotes the partition function in the torus  $\mathcal{T}_\varepsilon$ . Using (3.8) we write

$$\begin{aligned} & \sum_{\pi_\gamma \sigma_\gamma = s_\gamma} \exp[-\beta\gamma^{-d}H_\varepsilon(\sigma_\gamma)] \\ & \geq \sum_{\pi_\gamma \sigma_\gamma = s_\gamma} \exp[-\beta\gamma^{-d}H_\varepsilon(s_\gamma) - \beta c_2 \gamma^\delta (\gamma\varepsilon)^{-d}] \\ & = \exp[-\beta\gamma^{-d}H_\varepsilon(s_\gamma) - \beta c_2 \gamma^\delta (\gamma\varepsilon)^{-d}] W(s_\gamma) \end{aligned}$$

(and similarly for the upper bound). We call

$$\psi_\gamma = (\gamma\varepsilon)^{-d} \{c_2 \gamma^\delta + c_3 \gamma^{2(1-\delta)d} \log(\gamma^{-1})\} \quad (3.13)$$

and we get, using (3.9),

$$\begin{aligned} & \exp[-\gamma^{-d}[\beta H_\varepsilon(s_\gamma) - I_\varepsilon(s_\gamma)] - \psi_\gamma] \\ & \leq \sum_{\pi_\gamma \sigma_\gamma = s_\gamma} \exp[-\beta\gamma^{-d}H_\varepsilon(\sigma_\gamma)] \leq \exp[-\gamma^{-d}[\beta H_\varepsilon(s_\gamma) - I_\varepsilon(s_\gamma)] + \psi_\gamma] \end{aligned} \quad (3.14)$$

Calling  $m_{\beta,\gamma}$  the closest number to  $m_\beta$  which belongs to the range of  $s_\gamma(r)$ , we have

$$|m_\beta - m_{\beta,\gamma}| \leq c_5 \gamma^{(1-\delta)d} \quad (3.15)$$

and setting  $\hat{m}_{\beta,\gamma}(r) \equiv m_{\beta,\gamma}$ ,

$$|[\beta H_\varepsilon(\hat{m}_{\beta,\gamma}) - I_\varepsilon(\hat{m}_{\beta,\gamma})] - [\beta H_\varepsilon(\hat{m}_\beta) - I_\varepsilon(\hat{m}_\beta)]| \leq c_6 (\gamma\varepsilon)^{-d} \gamma^{(1-\delta)d}. \quad (3.16)$$

Therefore, using the lower bound in (3.14),

$$\begin{aligned} Z_{\beta,\gamma,\varepsilon} & \geq \sum_{\pi_\gamma \sigma_\gamma = \hat{m}_{\beta,\gamma}} \exp[-\beta\gamma^{-d}H_\varepsilon(\sigma_\gamma)] \\ & \geq \exp\{-\gamma^{-d}[\beta H_\varepsilon(\hat{m}_\beta) - I_\varepsilon(\hat{m}_\beta)] - \psi_\gamma - c_6 (\gamma\varepsilon)^{-d} \gamma^{(1-\delta)d}\} \end{aligned} \quad (3.17)$$

Hence, recalling (3.11), we find

$$\log(\mu_{\beta,\gamma,\varepsilon}(s_\gamma)) \leq -\gamma^{-d}\beta\mathcal{F}_\varepsilon(s_\gamma) + 2\psi_\gamma + c_6(\gamma\varepsilon)^{-d}\gamma^{(1-\delta)d} \quad (3.18)$$

For the upper bound we write

$$\begin{aligned} Z_{\beta,\gamma,\varepsilon} &= \sum_{\{s_\gamma\}} \sum_{\pi_\gamma \sigma_\gamma = s_\gamma} \exp[-\beta\gamma^{-d}H_\varepsilon(\sigma_\gamma)] \\ &\leq \sum_{\{s_\gamma\}} \exp\{-\gamma^{-d}[\beta H_\varepsilon(s_\gamma) - I_\varepsilon(s_\gamma)] + \psi_\gamma\} \end{aligned} \quad (3.19)$$

Since for all  $s_\gamma$

$$\beta H_\varepsilon(s_\gamma) - I_\varepsilon(s_\gamma) \geq \beta H_\varepsilon(\hat{m}_\beta) - I_\varepsilon(\hat{m}_\beta)$$

we get

$$\begin{aligned} \log Z_{\beta,\gamma,\varepsilon} &\leq -\gamma^{-d}[\beta H_\varepsilon(\hat{m}_\beta) - I_\varepsilon(\hat{m}_\beta)] + \psi_\gamma + \log N_\gamma \\ &\leq -\gamma^{-d}[\beta H_\varepsilon(\hat{m}_\beta) - I_\varepsilon(\hat{m}_\beta)] + \psi_\gamma + c_4(\gamma\varepsilon)^{-d}\gamma^{(1-\delta)d} \log(\gamma^{-1}) \end{aligned} \quad (3.20)$$

having used (3.10) in the last inequality. In conclusion

$$\log(\mu_{\beta,\gamma,\varepsilon}(s_\gamma)) \geq -\gamma^{-d}\beta\mathcal{F}_\varepsilon(s_\gamma) - 2\psi_\gamma - c_4(\gamma\varepsilon)^{-d}\gamma^{(1-\delta)d} \log(\gamma^{-1}) \quad (3.21)$$

The bounds (3.18) and (3.21) together with the obvious inequalities

$$\gamma^{(1-\delta)d} \log(\gamma^{-1}) \geq \gamma^{(1-\delta)d} \geq \gamma^{2(1-\delta)d} \log(\gamma^{-1})$$

prove (3.12). The proof of the Lemma is complete.  $\blacksquare$

By using Lemma 3.2 it is easy to prove Theorem 2.3:

*Proof of Theorem 2.3.* Recalling the definition (3.4) of  $\pi^{(k)}$ , for any  $\zeta^* > 0$  we set

$$\begin{aligned} \mathcal{A} &:= \{m \in L^\infty(\mathcal{T}_\varepsilon; [-1, 1]) : \\ &\quad \text{there is } r \in \mathbb{R}^d \text{ such that } |\pi^{(k)}m(r) - m_\beta| \geq \zeta^*\} \end{aligned} \quad (3.22)$$

$$\begin{aligned} \mathcal{B} &:= \{m \in L^\infty(\mathcal{T}_\varepsilon; [-1, 1]) \cap \mathcal{A}^c : \\ &\quad \text{there are } r', r'' \in \mathbb{R}^d \text{ such that } |\pi^{(k)}m(r') - m_\beta| < \zeta^* \\ &\quad \text{and } |\pi^{(k)}m(r'') + m_\beta| < \zeta^*\} \end{aligned} \quad (3.23)$$

We need the following lemma:

**Lemma.** For any  $k$  large enough and any  $\zeta^*$  small enough there is a constant  $c_7 > 0$  independent of  $\varepsilon$  such that

$$\inf_{m \in \mathcal{A} \cup \mathcal{B}} \mathcal{F}_\varepsilon(m) \geq c_7 \quad (3.24)$$

*Proof of the Lemma.* Suppose  $m \in \mathcal{A}$ . Call  $C = C^{(k)}(r)$  the cube in  $\mathcal{Q}^{(k)}$  that contains the point  $r$  appearing in (3.22). We set

$$\zeta := \frac{\zeta^*}{2}, \quad \alpha := \frac{\zeta|C|}{8}$$

Calling

$$\Lambda_\pm := \{r \in C : |m(r) \mp m_\beta| \leq \zeta\}; \quad \Lambda_0 := C \setminus (\Lambda_+ \cup \Lambda_-)$$

we first consider the case when  $|\Lambda_0| > \alpha$ . Then there is a function  $\kappa(\zeta) > 0$  such that

$$\mathcal{F}_\varepsilon(m) \geq \int_{\Lambda_0} dr [f(m(r)) - f(m_\beta)] \geq \kappa(\zeta)\alpha$$

When  $|\Lambda_0| \leq \alpha$  we may also suppose, without loss of generality, that  $|\Lambda_+| \geq |\Lambda_-|$ . By definition

$$\int_C dr m(r) < (m_\beta - \zeta^*)|C|$$

hence, since  $m \geq -1$  on  $\Lambda_0$ , we find

$$|\Lambda_+|(m_\beta - \zeta) - |\Lambda_0| + |\Lambda_-|(-m_\beta - \zeta) < (m_\beta - \zeta^*)|C|$$

Substituting  $|C| = |\Lambda_+| + |\Lambda_-| + |\Lambda_0|$ , we get

$$|\Lambda_-|(2m_\beta - \zeta^* + \zeta) > |\Lambda_+|(\zeta^* - \zeta) - 2|\Lambda_0|$$

Since  $|\Lambda_+| \geq (|C| - \alpha)/2$  we have

$$2|\Lambda_-| > \frac{|C| - \alpha}{2}\zeta - 2\alpha = \frac{|C|\zeta}{4} - \frac{\alpha\zeta}{2} \geq \frac{|C|\zeta}{8}$$

having supposed that  $m_\beta + \zeta \leq 1$ . Then

$$|\Lambda_+| \geq |\Lambda_-| \geq \frac{|C|\zeta}{16}$$



We can then conclude that in  $C$  there are two sets  $A_{\pm}$ ,  $|A_{\pm}| = |C|\zeta/16$ , where  $m(r)$  is respectively close to  $\pm m_{\beta}$  by at most  $\zeta$ ; moreover, by the isomorphism of Lebesgue measures, see [29], there is a one to one map  $\psi$  from  $A_+$  onto  $A_-$  which preserves the Lebesgue measure.

If  $|C|$  is small enough (i.e.  $k$  large), there is another cube  $C'$  in  $\mathcal{T}_{\varepsilon}$  and  $a > 0$  such that  $J(|r - r'|) \geq a$  for all  $r \in C$  and  $r' \in C'$ . We can then bound

$$\mathcal{F}_{\varepsilon}(m) \geq \frac{a}{4} \int_{C'} dr' \int_{A_+ \cup A_-} dr [m(r') - m(r)]^2$$

We write the integral over  $r$  as

$$\begin{aligned} & \int_{A_+} dr \{ [m(r') - m(r)]^2 + [m(r') - m(\psi(r))]^2 \} \\ & \geq \frac{1}{2} \int_{A_+} dr [m(\psi(r)) - m(r)]^2 \end{aligned}$$

thus proving the bound (3.24) limited to  $m \in \mathcal{A}$ .

If  $m \in \mathcal{B}$ , by definition it is not in  $\mathcal{A}$ , then, without loss of generality, we may suppose that the closures of the two cubes of  $\mathcal{Q}^{(k)}$  that contain  $r'$  and  $r''$ , see (3.23), have non empty intersection. We can then apply the same argument used for  $\mathcal{A}$  and the lemma is proved.  $\blacksquare$

We proceed in the proof of Theorem 2.3. Using (3.12), we have

$$\begin{aligned} & \mu_{\beta, \gamma, \varepsilon}(\{s^{(k)} \in \mathcal{A} \cup \mathcal{B}\}) \\ & \leq N_{\gamma} \exp\{-\beta c_7 \gamma^{-d} + (\gamma \varepsilon)^{-d} C_1 [\gamma^{\delta} + \gamma^{(1-\delta)d} \log(\gamma^{-1})]\} \end{aligned} \quad (3.25)$$

that vanishes as  $\gamma \rightarrow 0$  because of (3.10) and (3.3).

We have thus proved that the union of the two sets

$$\{|s^{(k)}(r) - m_{\beta}| < \zeta^*\}, \quad \{|s^{(k)}(r) + m_{\beta}| < \zeta^*\} \quad (3.26)$$

has probability going to 1 as  $\gamma \rightarrow 0^+$ , for any given  $k$ .

On the other hand, similarly to (3.7), we have, for any  $r \in \mathbb{R}^d$  and  $R > 0$ ,

$$\left| \int_{B_R(r)} dr' \sigma_{\gamma}(r') - \int_{B_R(r)} dr' \pi^{(k)} \sigma_{\gamma}(r') \right| \leq c_8 R^{-1} 2^{-k} \quad (3.27)$$

Therefore if  $s^{(k)}$  is in the first set in (3.26) then  $\sigma_{\gamma}$  is in the set  $\mathcal{P}_{R, \zeta, \gamma}^+$  of Definition 2.2 with  $\zeta = \zeta^* + c_8 R^{-1} 2^{-k}$ . Since the similar property holds

for  $\mathcal{P}_{R,\zeta,\gamma}^-$  we conclude that the statement in Theorem 2.3 is verified for the special case when  $\zeta_\gamma$  is equal to a constant independent of  $\gamma$ . Because this holds for any value of the constant, by a diagonalization procedure we conclude that there is a sequence  $\{\zeta_\gamma\}$  infinitesimal as  $\gamma \rightarrow 0^+$  for which the assertion remains true. The proof of Theorem 2.3 is thus completed.  $\blacksquare$

In the remaining part of this Section we will use Lemmas 3.1 and 3.2 to reduce the proof of Theorem 2.5 to the  $\Gamma$ -convergence problem for the functionals  $F_\varepsilon$  stated at the end of Section 2.

Let

$$m_\varepsilon^*(r) := u_\varepsilon^*(\varepsilon r), \quad r \in \mathcal{I}_\varepsilon \quad (3.28)$$

where  $\{u_\varepsilon^*\}$  is the minimizing sequence in the condition 2 for  $\Gamma$ -convergence. Let

$$\zeta_\gamma^* := \int_{\mathcal{I}} dr |u_\varepsilon^*(r) - u(r)| \quad (3.29)$$

(recall that  $\varepsilon^{-1} = \lceil \gamma^{-\alpha} \rceil$ ) and let  $R > 0$ ; we specify the sequence  $\zeta_\gamma$  in Theorem 2.5 as

$$\zeta_\gamma := \zeta_\gamma^* + c_1 R^{-1} \gamma^\delta \quad (3.30)$$

where  $c_1$  is defined in Lemma 3.1, see (3.7). The reason for this choice will become clear in the sequel.

Observe that given  $\zeta$  and  $R$  positive and setting

$$\zeta^\pm := \zeta \pm c_1 R^{-1} \gamma^\delta, \quad \text{thus } \zeta_\gamma^- = \zeta_\gamma^* \quad (3.31)$$

by (3.7) we have

$$\begin{aligned} \mu_{\beta,\gamma,\varepsilon}(\{\pi_\gamma \sigma_\gamma \in \mathcal{P}_{R,\zeta_\gamma^*,\gamma}(m)\}) &\leq \mu_{\beta,\gamma,\varepsilon}(\{\sigma_\gamma \in \mathcal{P}_{R,\zeta_\gamma,\gamma}(m)\}) \\ &\leq \mu_{\beta,\gamma,\varepsilon}(\{\pi_\gamma \sigma_\gamma \in \mathcal{P}_{R,\zeta_\gamma^+,\gamma}(m)\}) \end{aligned} \quad (3.32)$$

*The upper bound.* By (3.12)

$$\begin{aligned} &\log [\mu_{\beta,\gamma,\varepsilon}(\{\sigma_\gamma \in \mathcal{P}_{R,\zeta_\gamma,\gamma}(m)\})] \\ &\leq -\beta \gamma^{-d} \inf_{s_\gamma \in \mathcal{P}_{R,\zeta_\gamma^+,\gamma}(m)} \mathcal{F}_\varepsilon(s_\gamma) + \\ &\quad + \log N_\gamma + (\gamma \varepsilon)^{-d} C_1 \{\gamma^{(1-\delta)d} \log(\gamma^{-1}) + \gamma^\delta\} \end{aligned} \quad (3.33)$$

We multiply both sides by  $\gamma^d \varepsilon^{d-1}$  and let  $\gamma \rightarrow 0^+$ . By (3.10)

$$\lim_{\gamma \rightarrow 0^+} \gamma^d \varepsilon^{d-1} \log N_\gamma = 0$$

because  $\gamma^{(1-\delta)d\varepsilon^{-1}} = \gamma^{(1-\delta)d-\alpha}$  and by (3.2)  $(1-\delta)d > d\alpha \geq \alpha$ . Similarly also the last term in (3.33) vanishes, after having been multiplied by  $\gamma^d\varepsilon^{d-1}$ . Thus supposing the validity of the condition 1 of  $\Gamma$ -convergence, by (2.26) we get

$$\limsup_{\gamma \rightarrow 0^+} \gamma^d \varepsilon^{d-1} \log [\mu_{\beta, \gamma, \varepsilon}(\{\sigma_\gamma \in \mathcal{P}_{R, \zeta_\gamma, \gamma}(m)\})] \leq -\beta F(m) \quad (3.34)$$

*The lower bound.* We use (3.12), like in the upper bound, to get

$$\begin{aligned} & \log [\mu_{\beta, \gamma, \varepsilon}(\{\sigma_\gamma \in \mathcal{P}_{R, \zeta_\gamma, \gamma}(m)\})] \\ & \geq -\beta \gamma^{-d} \inf_{s_\gamma \in \mathcal{P}_{R, \zeta_\gamma^*, \gamma}(m)} \mathcal{F}_\varepsilon(s_\gamma) \\ & \quad - (\gamma\varepsilon)^{-d} C_1 \{\gamma^{(1-\delta)d} \log(\gamma^{-1}) + \gamma^\delta\} \end{aligned} \quad (3.35)$$

We have already seen in the proof of the upper bound that the last term in (3.35) multiplied by  $\gamma^d\varepsilon^{d-1}$  vanishes as  $\gamma \rightarrow 0^+$ . We would like to use condition 2 of the  $\Gamma$ -convergence, (2.27), for the first term on the right hand side of (3.35), but the minimizing sequence  $m_\varepsilon^*$  is not necessarily a coarse grained configuration. However, by (3.11), we have

$$\mathcal{F}_\varepsilon(m_\varepsilon^*) = [H_\varepsilon(m_\varepsilon^*) - I_\varepsilon(m_\varepsilon^*)] - [H_\varepsilon(\hat{m}_\beta) - I_\varepsilon(\hat{m}_\beta)] \quad (3.36)$$

Then, similarly to (3.8),

$$\gamma^{-d} |H_\varepsilon(m_\varepsilon^*) - H_\varepsilon(\pi_\gamma m_\varepsilon^*)| \leq c'_2 \gamma^\delta (\gamma\varepsilon)^{-d} \quad (3.37)$$

and, by the concavity of the entropy  $i(\cdot)$ , see (2.21),

$$-I_\varepsilon(m_\varepsilon^*) \geq -I_\varepsilon(\pi_\gamma m_\varepsilon^*) \quad (3.38)$$

$\pi_\gamma m_\varepsilon^*$  may not yet be a coarse grained function, but, recalling (3.15), there is a coarse grained function  $s_\gamma^*(r)$  such that

$$\sup_{r \in \mathcal{I}_\varepsilon} |\pi_\gamma m_\varepsilon^*(r) - s_\gamma^*(r)| \leq c'_5 \gamma^{(1-\delta)d} \quad (3.39)$$

Since the minimizing sequence  $\{m_\varepsilon^*\}$  has values in  $[-m_\beta, m_\beta]$  (see the end of Section 4) we have

$$|I_\varepsilon(\pi_\gamma m_\varepsilon^*) - I_\varepsilon(s_\gamma^*)| \leq c_9 (\gamma\varepsilon)^{-d} \gamma^{2(1-\delta)d} \quad (3.40)$$

By the continuity of the energy  $H_\varepsilon(\cdot)$  we then get

$$\gamma^{-d} \mathcal{F}_\varepsilon(m_\varepsilon^*) \geq \gamma^{-d} \mathcal{F}_\varepsilon(s_\gamma^*) - c'_2 \gamma^\delta (\gamma\varepsilon)^{-d} + c_{10} (\gamma\varepsilon)^{-d} \gamma^{2(1-\delta)d} \quad (3.41)$$

Thus, going back to the first term on the right hand side of (3.35),

$$\begin{aligned} & -\beta \gamma^{-d} \inf_{s_\gamma \in \mathcal{P}_{R, \zeta_\gamma^*, \gamma}(m)} \mathcal{F}_\varepsilon(s_\gamma) \\ & \geq -\beta \gamma^{-d} \mathcal{F}_\varepsilon(m_\varepsilon^*) - c'_2 \gamma^\delta (\gamma\varepsilon)^{-d} - c_{10} (\gamma\varepsilon)^{-d} \gamma^{2(1-\delta)d} \end{aligned} \quad (3.42)$$

We multiply both sides by  $\gamma^d \varepsilon^{d-1}$  and let  $\gamma \rightarrow 0^+$ . Then by (3.42) and (2.27),

$$\liminf_{\gamma \rightarrow 0^+} \log [\mu_{\beta, \gamma, \varepsilon}(\{\sigma_\gamma \in \mathcal{P}_{R, \zeta_\gamma, \gamma}(m)\})] \geq -\beta F(m) \quad (3.43)$$

(3.43) and (3.34) prove (2.16), hence Theorem 2.5 is reduced to the proof of (2.26) and (2.27), namely to the  $\Gamma$ -convergence of  $\{F_\varepsilon\}$  to  $F$ .

#### 4. GAMMA-CONVERGENCE

In this Section we prove that the functionals  $F_\varepsilon$  converge to  $F$  in the sense of  $\Gamma$ -convergence, see (2.26) and (2.27). While the proof of part 2 of  $\Gamma$ -convergence is standard the proof of part 1 is less typical, due to the non local structure of our functionals  $F_\varepsilon$ . As discussed at the end of Section 2, a local version of  $F_\varepsilon$  is the functional

$$M_\varepsilon(u) := \varepsilon \int_{\mathbb{R}^d} dr |\nabla u|^2 + \frac{1}{\varepsilon} \int_{\mathbb{R}^d} dr w(u)$$

where, for simplicity, we set  $w(s) := (1-s^2)^2$ . The proof of the  $\Gamma$ -convergence of  $\{M_\varepsilon\}$ , considered by Modica and Mortola, [26], and Modica, [25], exploits the following elementary inequality

$$M_\varepsilon(u) \geq 2 \int_{\mathbb{R}^d} dr |\nabla u| \sqrt{w(u)} =: L(u) \quad (4.1)$$

where  $\varepsilon$  has disappeared from the right hand side. Then the family  $\{L_\varepsilon\}$ ,  $L_\varepsilon \equiv L$ ,  $\Gamma$ -converges to the lower semicontinuous envelope of the functional  $L$ . On the other hand the minimizing sequence  $\{u_\varepsilon\}$  realizes equipartition of the energy, i.e.,

$$\varepsilon |\nabla u_\varepsilon|^2 = \frac{1}{\varepsilon} w(u_\varepsilon)$$

Hence  $M_\varepsilon(u_\varepsilon) = L(u_\varepsilon)$ .

In our case the inequality similar to (4.1) yields, for  $d = 1$ ,  $J = \mathbf{1}_{\{r \in \mathbb{R}: |r| \leq 1\}}/2$  and with  $[f(s) - f(m_\beta)]$  replaced by  $w(s)$ ,

$$F_\varepsilon(u) \geq \frac{1}{\varepsilon} \int_{\mathbb{R}} dr \left( \frac{1}{8\varepsilon} \int_{r-\varepsilon}^{r+\varepsilon} dr' |u(r) - u(r')|^2 \right)^{1/2} [w(u)]^{1/2} =: L_\varepsilon(u)$$

However the  $\Gamma$ -limit of  $\{L_\varepsilon\}$  (which corresponds in the previous case to the lower semicontinuous envelope of  $L$ ) vanishes. Indeed, let us consider  $u := 1$  on  $[0, 1]$  and  $u := -1$  elsewhere and choose  $u_\varepsilon := u$  for all  $\varepsilon$ . Then  $L_\varepsilon(u_\varepsilon) = 0$  for all  $\varepsilon$ , hence its limit vanishes.

Thus inequality (4.1) gives a trivial bound in our case and a different approach is required. The  $\Gamma$ -convergence of some non local functionals has also been considered recently by Jost, [21].

After these introductory remarks we begin the proofs. We need some notation. Recall that given a set  $E \subset \mathbb{R}^d$ ,  $\mathbf{1}_E$  denotes the characteristic function of  $E$ , i.e.  $\mathbf{1}_E(x) = 1$  if  $x \in E$  and  $\mathbf{1}_E(x) = 0$  otherwise. We call  $R = B \times [-h, h] \subset \mathcal{T}$  a parallelepiped in  $\mathcal{T}$  of height  $2h$  and middle section  $B$  (which is supposed in turn to be a parallelepiped in  $\mathbb{R}^{d-1}$  with  $2(d-1)$  faces). Thus  $B$  divides  $R$  into two parts that we call  $R^\pm$  and, according to this choice, we set  $\chi_R := m_\beta(\mathbf{1}_{R^+} - \mathbf{1}_{R^-})$ .

#### 4.1. Proof of condition 1 of Gamma-Convergence

**Theorem 4.1.** Let  $u \in BV(\mathcal{T}; \{\pm m_\beta\})$  and let  $\{u_\varepsilon\} \subset L^\infty(\mathcal{T}; [-1, 1])$  be a sequence converging to  $u$  in  $L^1(\mathcal{T})$  as  $\varepsilon \rightarrow 0^+$ . Then

$$\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) \geq s_\beta P(u)$$

*Proof.* Let  $u$  and  $u_\varepsilon$  be as in the statement of the Proposition. In Appendix D we shall prove that for any  $\zeta' > 0$  there are  $n$  disjoint parallelepipeds  $R_1, \dots, R_n$  with bases  $((d-1)$ -dimensional parallelepipeds)  $B_1, \dots, B_n$  respectively, and equal height  $2h$ , so that

$$\frac{1}{h} \sum_{i=1}^n \int_{R_i} dr |\chi_{R_i} - u| < \zeta', \quad \left| \sum_{i=1}^n |B_i| - P(u) \right| < \zeta' \quad (4.2)$$

By Proposition 4.2 below there exists an absolute constant  $c > 0$  such that for any  $\zeta > 0$

$$\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon; R_i) \geq (s_\beta - c\zeta) |B_i| - \frac{c}{h\zeta} \int_{R_i} dr |\chi_{R_i} - u|, \quad i = 1, \dots, n \quad (4.3)$$

By (2.24) we have  $F_\varepsilon(u_\varepsilon) \geq \sum_{i=1}^n F_\varepsilon(u_\varepsilon; R_i)$ . We then have, in view of (4.2),

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon) &\geq \sum_{i=1}^n \liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon; R_i) \\ &\geq (s_\beta - c\zeta) \sum_{i=1}^n |B_i| - \frac{c}{h\zeta} \sum_{i=1}^n \int_{R_i} dr |\chi_{R_i} - u| \\ &\geq s_\beta P(u) - c \frac{\zeta'}{\zeta} + O(\zeta) + O(\zeta') \end{aligned}$$

Taking  $\zeta' := \zeta^2$  and observing that the previous inequality holds for any  $\zeta > 0$ , the proof of the Proposition is concluded.  $\blacksquare$

**Proposition 4.2.** There is a constant  $c > 0$  so that the following holds. Let  $R \subset \mathcal{T}$  be a parallelepiped with basis  $B$  and height  $2h$ . If  $\{u_\varepsilon\} \subset L^\infty(\mathcal{T}; [-1, 1])$  converges in  $L^1(\mathcal{T})$  to  $u \in BV(\mathcal{T}; \{\pm m_\beta\})$ , then for any  $\zeta > 0$

$$\liminf_{\varepsilon \rightarrow 0^+} F_\varepsilon(u_\varepsilon; R) \geq (s_\beta - c\zeta)|B| - \frac{c}{h\zeta} \int_R dr |\chi_R - u| \quad (4.4)$$

Before proving this Proposition we introduce some definitions which shall be useful in the sequel.

Recalling Definition 2.1a we consider the partition  $\mathcal{Q}^{(-1)}$  of  $\mathbb{R}^d$  into cubes  $D$  of side 2 and denote by  $D(r)$  the cube  $D \in \mathcal{Q}^{(-1)}$  that contains the point  $r$ . Given  $D \in \mathcal{Q}^{(-1)}$ , we define the height  $\lambda(D)$  of  $D$  as  $\lambda(D) = 2n$ ,  $n \in \mathbb{Z}$ , if  $2n$  is the smallest value of the coordinate  $r_d$  of the points  $r \in D$ . Let  $R = B \times [-h, h]$ ,  $R_\varepsilon := \varepsilon^{-1}R \subset \mathcal{T}_\varepsilon := \varepsilon^{-1}\mathcal{T}$ ,  $B_\varepsilon := \varepsilon^{-1}B$ ,  $R_\varepsilon^\pm := \varepsilon^{-1}R^\pm$ . For simplicity we suppose that  $R_\varepsilon^\pm$  is  $\mathcal{Q}^{(-1)}$ -measurable, see Definition 2.1a.

For any  $m \in L^\infty(R_\varepsilon; [-m_\beta, m_\beta])$ , any positive integer  $k$  and  $\zeta > 0$ , we define the function  $\eta = \eta_{m,k,\zeta}$  as

$$\eta(r) := \begin{cases} 1 & \text{if } \pi^{(k)}m \geq m_\beta - \zeta \text{ on } D(r) \\ -1 & \text{if } \pi^{(k)}m \leq -m_\beta + \zeta \text{ on } D(r) \\ 0 & \text{otherwise} \end{cases} \quad (4.5)$$

where  $\pi^{(k)}$  is defined in (3.4).

We introduce now the important notion of cluster, see also Figure 1.

**Definition 4.3.** Given  $m \in L^\infty(\mathcal{T}_\varepsilon; [-1, 1])$  a cluster  $\mathcal{D} = \mathcal{D}(m)$  is a maximal  $*$ -connected union of cubes in  $R_\varepsilon^+$  where  $\eta(\cdot) < 1$  (two cubes are  $*$ -connected if their closures have non empty intersection).




Fig.1. The cubes  $D \in \mathcal{Q}^{(-1)} \cap R_\varepsilon^+$  where  $\eta < 1$ . There are three clusters: the one on the upper left is in  $\mathcal{L}^{(1)}$ , the other two in  $\mathcal{L}^{(0)}$ ; the one on the bottom left is in  $\mathcal{L}^{(0)} \setminus \mathcal{L}^*$  the other one in  $\mathcal{L}^*$ , as its height exceeds  $\varepsilon^{-1}h/2$ .




Fig.2. A cluster  $\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})$ .  $\mathcal{D}$  is divided into three parts, I, II, and III. The shadowed region II is the minimal section of  $\mathcal{D}$ . The modification  $m_\varepsilon^{(2)}$  of  $m_\varepsilon^{(1)}$  is equal everywhere to  $m_\varepsilon^{(1)}$  except in II, where it has value  $m_\beta$ ; thus  $\eta_{m_\varepsilon^{(2)},k,\zeta} = 1$  on II, hence III  $\in \mathcal{L}^{(1)}(m_\varepsilon^{(2)})$  and I  $\in \mathcal{L}^{(0)}(m_\varepsilon^{(2)})$  but I  $\notin \mathcal{L}^*(m_\varepsilon^{(2)})$ , which is thus empty.

$\mathcal{L}^0 = \mathcal{L}^0(m)$  is the set of clusters in  $R_\varepsilon^+$  which have non empty intersection with the basis  $B_\varepsilon$  of  $R_\varepsilon$ .  $\mathcal{L}^1 = \mathcal{L}^1(m)$  collects the others.

Given a cluster  $\mathcal{D}$ , we define its height  $\lambda(\mathcal{D})$  as

$$\lambda(\mathcal{D}) := \max\{\lambda(D) : D \in \mathcal{D}\}$$

and, for any even  $n$ , its section  $B(n; \mathcal{D})$  at height  $n$  as

$$B(n; \mathcal{D}) := \{D \subset \mathcal{D} : \lambda(D) = n\} \quad (4.6)$$

Calling

$$n^* := \frac{\varepsilon^{-1}h}{2} \quad (4.7)$$

(having supposed for simplicity that  $\varepsilon^{-1}h/2$  is even) we define

$$\mathcal{L}^* := \mathcal{L}^*(m) := \{\mathcal{D} \in \mathcal{L}^0 : \lambda(\mathcal{D}) \geq n^*\}$$

The minimal section  $S(\mathcal{D})$  of  $\mathcal{D} \in \mathcal{L}^*$ , see Figure 2, is defined after setting

$$\begin{aligned} b(\mathcal{D}) &:= \min_{\substack{n \text{ even} \\ 0 < n \leq n^*}} |B(n, \mathcal{D})| \\ n^0 &:= \min\{n \text{ even}, 0 < n \leq n^*, |B(n, \mathcal{D})| = b(\mathcal{D})\} \end{aligned}$$

as

$$S(\mathcal{D}) := B(n^0, \mathcal{D}) \quad (4.8)$$

Finally, the symbols  $c, c'$  will denote absolute positive constants that may change from line to line.

*Proof of Proposition 4.2.* Set

$$m_\varepsilon(r) := u_\varepsilon(\varepsilon r), \quad m(r) := u(\varepsilon r), \quad r \in \mathcal{F}_\varepsilon \quad (4.9)$$

We will prove the proposition after many successive modifications of the function  $m_\varepsilon$ , each one either decreasing  $\mathcal{F}_\varepsilon$  or increasing it by a “controlled” quantity. The first step is very simple, we just take

$$m_\varepsilon^{(1)}(r) := \begin{cases} \pm m_\beta & \text{if } m_\varepsilon(r) \geq m_\beta, \text{ respectively } m_\varepsilon(r) \leq -m_\beta \\ m_\varepsilon(r) & \text{otherwise} \end{cases}$$

Then, obviously,  $\mathcal{F}_\varepsilon(m_\varepsilon) \geq \mathcal{F}_\varepsilon(m_\varepsilon^{(1)})$ .

Modulo rotation and reflection of the axes, we may suppose that the basis of  $R_\varepsilon$  is contained in the coordinate hyperplane  $\{r_d = 0\}$ ,  $r_d$  denoting the last coordinate of  $r$  in  $\mathbb{R}^d$ , and that  $R_\varepsilon^+ \subset \{r_d > 0\}$ .

We next modify  $m_\varepsilon^{(1)}$  in  $R_\varepsilon^+$  (the modification in  $R_\varepsilon^-$  is similar and done later) by cutting the clusters in  $\mathcal{L}^*(m_\varepsilon^{(1)})$  at their minimal section, see Figure



2. The clusters are now defined by means of the function  $\eta_{m_\varepsilon^{(1)},k,\zeta}$ , see (4.5). Let

$$m_\varepsilon^{(2)} := \begin{cases} m_\beta & \text{on } S(\mathcal{D}), \mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)}) \\ m_\varepsilon^{(1)} & \text{elsewhere on } R_\varepsilon \end{cases}$$

Then

$$\mathcal{F}(m_\varepsilon^{(1)}; R_\varepsilon) \geq \mathcal{F}(m_\varepsilon^{(2)}; R_\varepsilon) - c \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \quad (4.10)$$

We shall see at the end of the proof of this proposition, that the cost of this substitution, that is the last term in (4.10), can be controlled in terms of the  $L^1(R)$  norm of  $\chi_R - u_\varepsilon$ .

By construction  $\mathcal{L}^*(m_\varepsilon^{(2)}) = \emptyset$ , so that the “dangerous clusters” that intersect the basis of  $R_\varepsilon$ , i.e., those in  $\mathcal{L}^0(m_\varepsilon^{(2)})$ , are all “low”, in the sense that they do not reach the height  $n^*$ , which is half of the total height of  $R_\varepsilon^+$ . We next apply Theorem B.2 of Appendix B to modify  $m_\varepsilon^{(2)}$  into a function  $m_\varepsilon^{(3)}$ , which, as we shall see, is positive on the clusters of  $\mathcal{L}^1(m_\varepsilon^{(2)})$ , where instead  $m_\varepsilon^{(2)}$  may be negative: the proof exploits the fact that by definition all the clusters of  $\mathcal{L}^1(m_\varepsilon^{(2)})$  are surrounded by cubes where  $\eta(\cdot) = 1$ . The precise statement can be found in Theorem B.2 that we apply in the present context with  $\Delta$ ,  $\Gamma$  and  $\Lambda$  defined as follows:

$$\begin{aligned} \Delta &:= \cup \{ \mathcal{D} : \mathcal{D} \in \mathcal{L}^1(m_\varepsilon^{(2)}) \} \\ \Gamma &:= \{ D \in \mathcal{Q}^{(-1)} \cap R_\varepsilon^+ : D \cap \Delta = \emptyset, \bar{D} \cap \Delta \neq \emptyset \} \\ \Lambda &:= R_\varepsilon \setminus (\Delta \cup \Gamma) \end{aligned}$$

The parameters  $\zeta$  and  $k$  of Theorem B.2 are the same parameters as in the definition of the functions  $\eta$  in (4.5);  $\ell$  is a number in  $(0, 1)$  and the sequence  $\{c_k\}$  (defined in (B.7)) is actually independent of  $\varepsilon$  (see the Remark at the end of the proof of Lemma B.1) and tends to 0 as  $k \rightarrow +\infty$ ; finally we recall that  $0 < \theta := \zeta + c_k < m_\beta$ . We choose  $\zeta$  and  $k$  so that  $\theta = \zeta + c_k \leq m_\beta/4$ .

Since  $m_\varepsilon^{(2)}$  satisfies condition (B.4), i.e.,  $\pi^{(k)} m_\varepsilon^{(2)} \geq m_\beta - \zeta$  for all  $r \in \Gamma$ , then, by Theorem B.2, there exists  $m_\varepsilon^{(3)} := (m_\varepsilon^{(2)})_{\ell, \theta}$  on  $R_\varepsilon$  with the following properties:

1.  $\mathcal{F}(m_\varepsilon^{(2)}; R_\varepsilon) \geq \mathcal{F}(m_\varepsilon^{(3)}; R_\varepsilon)$ ;
2.  $m_\varepsilon^{(3)} \geq m_\varepsilon^{(2)}$  on  $R_\varepsilon$ , and  $m_\varepsilon^{(3)}(r) = m_\varepsilon^{(2)}(r)$  for all  $r \in R_\varepsilon$  at distance not smaller than 1 from  $\Delta$ ;
3.  $m_\varepsilon^{(3)} \geq m_\beta - \theta$  on  $\Delta$ .




Fig.3.  $\mathcal{L}^{(1)}(m_\varepsilon^{(3)}) = \mathcal{L}^*(m_\varepsilon^{(3)}) = \emptyset$ , therefore  $\eta = 1$  on  $\{\lambda(D) \geq n^* + 4\}$ . In particular  $\eta = 1$  on the level  $\lambda(D) = n^* + 4$ , which is shadowed. While the averages  $\pi_k m_\varepsilon^{(3)} \geq m_\beta - \theta$  above this level, this may not be so point by point. However for the modification  $m_\varepsilon^{(4)}$  of  $m_\varepsilon^{(3)}$ ,  $m_\varepsilon^{(4)} \geq m_\beta - \theta$  on  $\{\lambda(D) > n^* + 4\}$ .

By construction  $-m_\beta \leq m_\varepsilon^{(3)} \leq m_\beta$ ,  $\mathcal{L}^*(m_\varepsilon^{(3)}) = \emptyset$ , and  $m_\varepsilon^{(3)} \geq m_\beta - \theta$  on all clusters  $\mathcal{D}$  with  $\mathcal{D} \cap B_\varepsilon = \emptyset$  where  $\eta_{m_\varepsilon^{(2)}, k, \zeta} < 1$  (i.e., where  $m_\varepsilon^{(2)}$  is “far” from the value  $m_\beta$ ).

We redefine the clusters (see (4.5)) for  $m_\varepsilon^{(3)}$  with  $\zeta$  replaced by  $\theta$ , i.e., by means of the function  $\eta_{m_\varepsilon^{(3)}, k, \theta}$ . By an abuse of notation the new clusters are denoted by the same symbols as the old ones and we also denote  $\theta$  again by  $\zeta$ . By construction, if  $\mathcal{D} \in \mathcal{L}^0(m_\varepsilon^{(3)})$ , then  $B(n, \mathcal{D}) = \emptyset$  for  $n > n^*$ , see (4.6) for notation.

We now modify  $m_\varepsilon^{(3)}$  into  $m_\varepsilon^{(4)}$  in such a way that  $m_\varepsilon^{(4)} \geq m_\beta - \theta$  on all  $D \in \mathcal{Q}^{(-1)}$  with  $\lambda(D) \geq n^* + 4$ , see Figure 3. Precisely we apply again Theorem B.2 with  $\Delta$ ,  $\Gamma$  and  $\Lambda$  defined as follows:

$$\begin{aligned} \Delta &:= \{D \in \mathcal{Q}^{(-1)} \cap R_\varepsilon^+ : \lambda(D) \geq n^* + 4\} \\ \Gamma &:= \{D \in \mathcal{Q}^{(-1)} \cap R_\varepsilon^+ : \lambda(D) = n^* + 2\} \\ \Lambda &:= R_\varepsilon \setminus (\Delta \cup \Gamma) \end{aligned}$$

(note that  $\eta_{m_\varepsilon^{(2)}, k, \zeta}(\cdot) = 1$  on  $\Gamma$  so that  $m_\varepsilon^{(2)}$  satisfies condition (B.4)). Then by Theorem B.2 there exists  $m_\varepsilon^{(4)}$  on  $R_\varepsilon$  with the following properties:

1.  $\mathcal{F}(m_\varepsilon^{(3)}; R_\varepsilon) \geq \mathcal{F}(m_\varepsilon^{(4)}; R_\varepsilon)$ ;
2.  $m_\varepsilon^{(4)} \geq m_\varepsilon^{(3)}$  on  $R_\varepsilon$ , and  $m_\varepsilon^{(4)} = m_\varepsilon^{(3)}$  on all  $D$  with  $\lambda(D) < n^* + 2$ ;

3.  $m_\varepsilon^{(4)} \geq m_\beta - \theta$  on  $\Delta$ , i.e., on all  $D$  with  $\lambda(D) \geq n^* + 4$ .

We conclude this first part of the proof of Proposition 4.2 by introducing the function  $m_\varepsilon^{(5)}$ , obtained by repeating (with opposite sign) the modifications that led to  $m_\varepsilon^{(4)}$  also in the lower part  $R_\varepsilon^-$  of the parallelepiped  $R_\varepsilon$ .

**Summary of What Has Been Done so Far.** There is a function  $m_\varepsilon^{(5)}$  which is larger than  $m_\beta - \theta$  for  $r_d \geq n^* + 4$  and smaller than  $-m_\beta + \theta$  for  $r_d \leq -n^* - 4$ , such that

$$\mathcal{F}(m_\varepsilon; R_\varepsilon) \geq \mathcal{F}(m_\varepsilon^{(5)}; R_\varepsilon) - c \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \quad (4.11)$$

(here  $\mathcal{L}^*$  is referred to the whole  $R_\varepsilon$ ).

Let  $K_\varepsilon := B_\varepsilon \times (-\infty, +\infty)$ ,

$$m_\varepsilon^{(6)} := \begin{cases} m_\varepsilon^{(5)} & \text{on } R_\varepsilon \\ \pm m_\beta & \text{on } K_\varepsilon^\pm \setminus R_\varepsilon^\pm \end{cases}$$

Then one can verify that

$$\mathcal{F}(m_\varepsilon^{(5)}; R_\varepsilon) \geq \mathcal{F}(m_\varepsilon^{(6)}; K_\varepsilon) - c\zeta|B_\varepsilon| \quad (4.12)$$

(choosing  $k$  so large that  $c_k < \zeta$ ).

We would like to have  $\mathcal{F}(m_\varepsilon^{(6)}; K_\varepsilon) \geq \mathcal{F}(m^*; K_\varepsilon)$  when  $m^*$  is the instanton  $\bar{m}$  of Theorem 2.6 on each line parallel to the  $r_d$ -axis of  $K_\varepsilon$ . However we are not able to make an estimate of  $\mathcal{F}(m_\varepsilon^{(6)}; K_\varepsilon)$  to verify the inequality. A possible way out would be to prove that the infimum of  $\mathcal{F}(\cdot; K_\varepsilon)$  over all  $m \in L^\infty(K_\varepsilon; [-m_\beta, m_\beta])$  having asymptotic values  $\pm m_\beta$  for  $r_d \rightarrow \pm\infty$  is just  $\mathcal{F}(m^*; K_\varepsilon)$ . This is however ruled out by the fact that  $m^*$  is not a critical point of  $\mathcal{F}(\cdot; K_\varepsilon)$ . In fact, all critical points of  $\mathcal{F}(\cdot; K_\varepsilon)$  (see the Remark below) must verify

$$m(r) = \tanh\{\beta \tilde{J} * m(r)\}, \quad \tilde{J} * m(r) := \int_{K_\varepsilon} dr' \tilde{J}(r, r') m(r') \quad (4.13)$$

where

$$\begin{aligned} \tilde{J}(r, r') &:= J(|r - r'|) \mathbf{1}_{\{r, r' \in K_\varepsilon\}} + j(r) \delta(r - r') \\ j(r) &:= \int_{\{r'' \notin K_\varepsilon\}} dr'' J(|r - r''|) \end{aligned} \quad (4.14)$$

and  $\delta(r - r')$  is the delta function. Since  $m^*(r) = \tanh\{\beta J * m^*(r)\}$  when the convolution is over the whole  $\mathbb{R}^d$ , in general  $m^*(r) \neq \tanh\{\beta \tilde{J} * m^*(r)\}$  when the distance of  $r$  from  $\partial K_\varepsilon$  is smaller than 1.

**Remark.** The Euler-Lagrange equation for the functional  $\mathcal{F}(m) := \mathcal{F}(m; \mathbb{R}^d)$  is

$$f'(m) + m = J * m \quad (4.15)$$

Recalling the definition of  $f$  in (2.20), (4.15) becomes

$$-i'(m) = \beta J * m$$

Since  $-i'(s) = \operatorname{arctanh}(s)$ , one deduces that a function  $m \in L^\infty(\mathbb{R}^d; (-1, 1))$  solves (4.15) if and only if

$$m = \tanh(\beta J * m)$$

To overcome the problems due to the presence of the delta function in the interaction, we introduce an auxiliary functional  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon) \leq \mathcal{F}(\cdot; K_\varepsilon)$  whose critical points can be characterized. We will see that there is a unique critical point which is an instanton function close to  $m^*$ . The minimum is found by means of an auxiliary dynamics on  $L^\infty(K_\varepsilon; [-1, 1])$  under which the functional  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon)$  is monotonic (non-increasing). The minimum is then obtained as the limit point when  $t \rightarrow +\infty$  of the orbit  $m_t$  that starts from  $m_\varepsilon^{(6)}$ . The analysis adapts to the present context results known in the literature, [13] and [14]. Even though the main ideas are the same, the extension is not totally trivial and we will report some details in Appendix C. The dynamics here is merely a technical tool, but the evolution has an intrinsic interest with many significant applications in the physics of phase separation and mathematical interest in its own right, see for instance [9], [10], [11], [12].

As explained above a way to avoid the delta function in (4.14) is to modify  $\mathcal{F}(\cdot; K_\varepsilon)$  into another functional,  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon)$ , which does not produce anymore the dangerous local term in the interaction. For any  $m \in L^\infty(K_\varepsilon; [-1, 1])$  we set

$$\begin{aligned} \mathcal{F}^{(1)}(m; K_\varepsilon) &:= \int_{K_\varepsilon} dr (1 - j(r)) [f(m(r)) - f(m_\beta)] \\ &+ \frac{1}{4} \iint_{K_\varepsilon \times K_\varepsilon} dr dr' J(|r - r'|) [m(r) - m(r')]^2 \end{aligned} \quad (4.16)$$

Obviously

$$\mathcal{F}^{(1)}(m; K_\varepsilon) \leq \mathcal{F}(m; K_\varepsilon) \quad (4.17)$$

and since

$$\int_{K_\varepsilon} dr (1 - j(r)) m(r)^2 = \iint_{K_\varepsilon \times K_\varepsilon} dr dr' m(r)^2 J(|r - r'|)$$

we have, recalling (2.20),

$$\begin{aligned} \mathcal{F}^{(1)}(m; K_\varepsilon) &= -\beta^{-1} \int_{K_\varepsilon} dr [1 - j(r)] i(m(r)) \\ &\quad - \frac{1}{2} \iint_{K_\varepsilon \times K_\varepsilon} dr dr' J(|r - r'|) m(r) m(r') - C^{(1)}(\beta) \end{aligned} \quad (4.18)$$

where  $C^{(1)}(\beta)$  is the sum of the first two terms when  $m(\cdot) \equiv m_\beta$ .

By direct inspection  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon)$  does not increase along the solutions of the equation

$$\begin{aligned} \frac{dm_t(r)}{dt} &= -m_t(r) + \tanh\{\beta J^{(1)} * m_t(r)\} \\ J^{(1)} * m_t(r) &:= \int_{K_\varepsilon} dr' J^{(1)}(r, r') m_t(r') \end{aligned} \quad (4.19)$$

where

$$J^{(1)}(r, r') := \frac{1}{1 - j(r)} J(|r - r'|), \quad \int_{K_\varepsilon} dr' J^{(1)}(r, r') = 1 \quad (4.20)$$

We show in Appendix C that there is a (instanton-like) function  $m_\varepsilon^{(7)}$  on  $K_\varepsilon$ , which is a stationary solution of (4.19) in the whole  $K_\varepsilon$ . This solution is an antisymmetric function of  $r_d$  and there are  $c > 0$  and  $c'$  independent of  $\varepsilon$  and of the section  $B_\varepsilon$  of the cylinder  $K_\varepsilon$ , so that

$$m_\varepsilon^{(7)}(r) \geq m_\beta - c' e^{-cr_d}, \quad r_d \geq 0 \quad (4.21)$$

Moreover, if  $m_t$  is the orbit starting from  $m_\varepsilon^{(6)}$ , then there is  $r^0 = (0, r_d^0)$  so that

$$\lim_{t \rightarrow +\infty} m_t(r) = m_\varepsilon^{(7)}(r - r^0) \quad (4.22)$$

uniformly on the compact subsets of  $K_\varepsilon$ . By the lower semicontinuity of  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon)$  and its invariance under vertical translations,

$$\mathcal{F}^{(1)}(m_\varepsilon^{(6)}; K_\varepsilon) \geq \liminf_{t \rightarrow +\infty} \mathcal{F}^{(1)}((m_\varepsilon^{(6)})_t; K_\varepsilon) \geq \mathcal{F}^{(1)}(m_\varepsilon^{(7)}; K_\varepsilon) \quad (4.23)$$

By (4.11), (4.12), (4.17) and (4.23) we have

$$\begin{aligned} \mathcal{F}(m_\varepsilon; R_\varepsilon) &\geq \mathcal{F}^{(1)}(m_\varepsilon^{(6)}; K_\varepsilon) - c\zeta|B_\varepsilon| - c \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \\ &\geq \mathcal{F}^{(1)}(m_\varepsilon^{(7)}; K_\varepsilon) - c\zeta|B_\varepsilon| - c \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \end{aligned} \quad (4.24)$$

where  $\mathcal{L}^*$  is referred to the whole  $R_\varepsilon$ .

Our last effort will be to replace  $m_\varepsilon^{(7)}$  by  $m^*$  which we recall is the instanton  $\tilde{m}$  of Theorem 2.6 on each line parallel to the  $r_d$ -axis of  $K_\varepsilon$ , with the 0's belonging to the original middle section  $B_\varepsilon$  of  $R_\varepsilon$ . The natural way would be to prove that  $m_\varepsilon^{(7)}$  is really close to  $m^*$  except possibly at points close to the boundaries and with height  $r_d$  not too large. Recall in fact that both  $m_\varepsilon^{(7)}$  and  $m^*$  have the same asymptotic values that they approach exponentially fast. However, although we believe this statement to be correct, we miss a proof. We will proceed by changing the functional and the corresponding dynamics with the introduction of “Neumann conditions” at the boundaries. After that, the above comparison will become easier. To construct the kernel  $J^{(2)}$  we need to smooth the “corners” of  $B_\varepsilon$  from inside; hence let  $\tilde{B}_\varepsilon \subset B_\varepsilon$  be a convex  $C^\infty$  set (in Appendix D it is shown that  $B$  can be taken as a cube) with

$$|B_\varepsilon \setminus \tilde{B}_\varepsilon| \leq c\varepsilon^{2-d}$$

Let  $\tilde{K}_\varepsilon := \tilde{B}_\varepsilon \times (-\infty, +\infty) \subset K_\varepsilon$ . Then, denoting again by  $m_\varepsilon^{(7)}$  the restriction of this function to  $\tilde{K}_\varepsilon$ , we have

$$\mathcal{F}^{(1)}(m_\varepsilon^{(7)}; K_\varepsilon) \geq \mathcal{F}^{(1)}(m_\varepsilon^{(7)}; \tilde{K}_\varepsilon) - c\varepsilon^{2-d} \quad (4.25)$$

because of the exponential convergence of  $m_\varepsilon^{(7)}$  to  $\pm m_\beta$  as  $r_d \rightarrow \pm\infty$ .

We denote by  $d(r, \partial\tilde{K}_\varepsilon)$  the distance of  $r \in \tilde{K}_\varepsilon$  from the boundary  $\partial\tilde{K}_\varepsilon$  of  $\tilde{K}_\varepsilon$ . In Appendix C we prove that there is a smooth function  $J^{(2)}(r, r') : \tilde{K}_\varepsilon \times \tilde{K}_\varepsilon \rightarrow [0, 1]$  with the following properties:

1.  $J^{(2)}$  is supported on  $\{(r, r') : r \in \tilde{K}_\varepsilon, r' \in B_1(r) \cap \tilde{K}_\varepsilon\}$ ,
2.  $J^{(2)}(r, r') = J^{(2)}(r', r)$  on  $\tilde{K}_\varepsilon \times \tilde{K}_\varepsilon$ ,
3.  $\int_{\tilde{K}_\varepsilon} dr' J^{(2)}(r, r') = 1$  for all  $r \in \tilde{K}_\varepsilon$ ,
4.  $J^{(2)}(r, r') = J(|r - r'|)$  for all  $r, r' \in \tilde{K}_\varepsilon$  such that  $d(r, \partial\tilde{K}_\varepsilon) \geq 1$ ,
5. set  $r = (\xi, r_d)$ ,  $\xi \in \tilde{B}_\varepsilon$ ,  $r_d \in \mathbb{R}$ ; then for all  $\xi \in \tilde{B}_\varepsilon$ , and all  $r_d, r'_d \in \mathbb{R}$

$$\int_{\tilde{B}_\varepsilon} d\xi' J^{(2)}((\xi, r_d), (\xi', r'_d)) = \int_{\tilde{B}_\varepsilon} d\xi' J(|(\xi, r_d) - (\xi', r'_d)|) \quad (4.26)$$

We then define for any  $m \in L^\infty(\tilde{K}_\varepsilon; [-1, 1])$

$$\begin{aligned} \mathcal{F}^{(2)}(m; \tilde{K}_\varepsilon) &:= \int_{\tilde{K}_\varepsilon} dr [f(m(r)) - f(m_\beta)] \\ &\quad + \frac{1}{4} \iint_{\tilde{K}_\varepsilon \times \tilde{K}_\varepsilon} dr dr' J^{(2)}(r, r') [m(r) - m(r')]^2 \end{aligned} \quad (4.27)$$

The same argument used to prove (4.25) shows that

$$\mathcal{F}^{(1)}(m_\varepsilon^{(7)}; \tilde{K}_\varepsilon) \geq \mathcal{F}^{(2)}(m_\varepsilon^{(7)}; \tilde{K}_\varepsilon) - c\varepsilon^{2-d} \quad (4.28)$$

By direct inspection one can prove that  $\mathcal{F}^{(2)}(m_t; \tilde{K}_\varepsilon)$  is non-increasing along any orbit  $m_t$  solution of (4.19) with  $K_\varepsilon$  replaced by  $\tilde{K}_\varepsilon$  and  $J^{(1)}(r, r')$  by  $J^{(2)}(r, r')$ .

Let  $\psi_t \in L^\infty(\mathbb{R}; [-1, 1])$ ,  $t \geq 0$ , satisfy the equation

$$\frac{d\psi_t(s)}{dt} = -\psi_t(s) + \tanh\{\beta \bar{J} * \psi_t(s)\}, \quad s \in \mathbb{R} \quad (4.29)$$

with  $\bar{J}$  as in (2.18). Then by statement 5 the function  $m_t \in L^\infty(\tilde{K}_\varepsilon; [-1, 1])$ ,  $t \geq 0$ , defined by  $m_t(r) := \psi_t(r_d)$  solves (4.19) with  $\tilde{K}_\varepsilon$  and  $J^{(2)}$ . As a consequence,  $m^*(r) = \bar{m}(r_d)$ ,  $r \in \tilde{K}_\varepsilon$ ,  $\bar{m}$  as in Theorem 2.6, is a stationary solution of this new version of (4.19).

Let  $m_t \in L^\infty(\tilde{K}_\varepsilon; [-1, 1])$  solve this new version of (4.19) with initial condition  $m_0 = m_\varepsilon^{(7)}$ . Then, by arguments completely similar to the previous ones which led to  $m_\varepsilon^{(7)}$ , we conclude that  $m_t \rightarrow m^*$  as  $t \rightarrow +\infty$ . Thus

$$\mathcal{F}^{(2)}(m_\varepsilon^{(7)}; \tilde{K}_\varepsilon) \geq \mathcal{F}^{(2)}(m^*; \tilde{K}_\varepsilon) \quad (4.30)$$

Finally, there is a constant  $c > 0$  so that, denoting by  $|B|$  the area of the original basis  $B$  of  $R$ ,

$$\mathcal{F}^{(2)}(m^*; \tilde{K}_\varepsilon) = \varepsilon^{1-d} s_\beta |B| - c\varepsilon^{2-d} \quad (4.31)$$

By (4.24), (4.28), (4.30) and (4.31) we have

$$\begin{aligned} \mathcal{F}(m_\varepsilon; R_\varepsilon) &\geq \mathcal{F}^{(2)}(m^*; \tilde{K}_\varepsilon) - c\varepsilon^{2-d} - c\zeta |B_\varepsilon| - c \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \\ &\geq \varepsilon^{1-d} (s_\beta - c\zeta) |B| - c\varepsilon^{2-d} - c \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \end{aligned} \quad (4.32)$$

where  $\mathcal{L}^*$  is referred to the whole  $R_\varepsilon$ . We first consider the sum relative to  $R_\varepsilon^+$ . By the definition of minimal section, if  $\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})$

$$|S(\mathcal{D})|n^* \leq |\mathcal{D}| \quad (4.33)$$

with  $n^*$  as in (4.7). Moreover,  $\eta_{m_\varepsilon^{(1)},k,\zeta} < 1$  on any  $D \subset \mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})$ , so that in any such  $D$  there is a cube  $Q \in \mathcal{Q}^{(k)}$  of side  $2^{-k}$  where

$$\frac{1}{|Q|} \int_Q dr m_\varepsilon^{(1)}(r) \leq m_\beta - \zeta \quad (4.34)$$

Thus,

$$\begin{aligned} \int_D dr |\chi_R(\varepsilon r) - m_\varepsilon^{(1)}(r)| &\geq \int_Q dr |\chi_R(\varepsilon r) - m_\varepsilon^{(1)}(r)| \\ &= \int_Q dr [m_\beta - m_\varepsilon^{(1)}(r)] \\ &\geq |Q|\zeta = 2^{-(k+1)d}|D|\zeta \end{aligned} \quad (4.35)$$

because  $Q \subset R_\varepsilon^+$  where  $\chi_R = m_\beta$ . By (2.23),  $\mathcal{F}_\varepsilon = \varepsilon^{1-d}F_\varepsilon$ , hence, using (4.33) and recalling (4.7) where  $n^*$  is defined, and (4.9), we have

$$\begin{aligned} &\varepsilon^{d-1} \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} |S(\mathcal{D})| \\ &\leq \varepsilon^d \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} \frac{2|\mathcal{D}|}{h} \\ &\leq \varepsilon^d \sum_{\mathcal{D} \in \mathcal{L}^*(m_\varepsilon^{(1)})} \sum_{D \subset \mathcal{D}} \frac{2^{(k+2)d}}{h\zeta} \int_D dr |\chi_R(\varepsilon r) - m_\varepsilon^{(1)}(r)| \\ &\leq \frac{c}{h\zeta} \int_R dr |\chi_R(r) - u_\varepsilon^{(1)}(r)| \\ &\leq \frac{c}{h\zeta} \int_R dr |\chi_R(r) - u_\varepsilon(r)| \end{aligned} \quad (4.36)$$

where  $u_\varepsilon^{(1)}(r) := m_\varepsilon^{(1)}(\varepsilon^{-1}r)$ ,  $r \in \mathcal{T}$ . The same bound is obtained when we sum over the clusters in  $R_\varepsilon^-$ . By (4.32) and (4.36) we then deduce that

$$F_\varepsilon(u_\varepsilon; R) = \varepsilon^{d-1} \mathcal{F}(m_\varepsilon; R_\varepsilon) \geq (s_\beta - c\zeta)|B| - c\varepsilon - \frac{c}{h\zeta} \int_R dr |\chi_R(r) - u_\varepsilon(r)|$$



Then recalling that  $u_\varepsilon \rightarrow u$  in  $L^1(\mathcal{I})$  as  $\varepsilon \rightarrow 0^+$ , we obtain (4.4). The Proposition is thus proved.  $\blacksquare$

## 4.2. Proof of condition 2 of Gamma-Convergence

We only sketch the proof. Let  $u \in BV(\mathcal{I}; \{\pm m_\beta\})$ . We first suppose that the boundary  $\partial E$  of the set  $\{u(r) = m_\beta\}$  is a hypersurface of class  $\mathcal{C}^1$ . Let  $m(r) := u(\varepsilon r)$ ,  $r \in \mathcal{I}_\varepsilon$ , so that  $m \in L^\infty(\mathcal{I}_\varepsilon; \{\pm m_\beta\})$ . Given  $0 < \delta < 1$  and  $\varepsilon > 0$ , we define  $u_\varepsilon \in L^\infty(\mathcal{I}_\varepsilon; [-m_\beta, m_\beta])$  as follows. Let  $d(r)$  be the signed distance function from  $\varepsilon^{-1}\partial E$  positive inside  $\varepsilon^{-1}E$ . We then set

$$m_\varepsilon^*(r) := \begin{cases} \bar{m}(d(r)) & \text{if } |d(r)| \leq \varepsilon^{-\delta} \\ m_\beta & \text{if } d(r) > \varepsilon^{-\delta} \\ -m_\beta & \text{if } d(r) < -\varepsilon^{-\delta} \end{cases}$$

where  $\bar{m}(s)$  is the instanton of Theorem 2.6 which converges to  $\pm m_\beta$  as  $s \rightarrow \pm\infty$  exponentially fast. We then have, using the coarea formula (see 3.4.4 in [17])

$$\begin{aligned} \mathcal{F}_\varepsilon(m_\varepsilon^*) &\leq c'e^{-c\varepsilon^{-\delta}} + \int_{-\varepsilon^{-\delta}}^{\varepsilon^{-\delta}} dt [f(\bar{m}(t)) - f(m_\beta)] \lambda(\{d(x) = t\}) + \\ &+ \frac{1}{4} \int_{-\varepsilon^{-\delta}}^{\varepsilon^{-\delta}} dt \int_{-\varepsilon^{-\delta}}^{\varepsilon^{-\delta}} ds [\bar{m}(t) - \bar{m}(s)]^2 \int_{\{d(x)=t\}} d\lambda(x) \times \\ &\times \int_{\{d(y)=s\}} d\lambda(y) J(|(x,t) - (y,s)|) \end{aligned}$$

where  $d\lambda$  denotes the  $(d-1)$ -dimensional surface measure. Now for any  $(x, t)$  as above

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\{d(y)=s\}} d\lambda(y) J(|(x,t) - (y,s)|) = \bar{J}(|t-s|)$$

see (2.18), and

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon^{d-1} \int_{\{d(x)=t\}} d\lambda(x) = |\partial E|$$

Hence

$$\limsup_{\varepsilon \rightarrow 0^+} \varepsilon^{d-1} \mathcal{F}_\varepsilon(m_\varepsilon^*) \leq \bar{\mathcal{F}}(\bar{m})|\partial E| = s_\beta|\partial E|$$

Setting  $u_\varepsilon^*(r) := m_\varepsilon^*(\varepsilon^{-1}r)$ ,  $r \in \mathcal{T}$ , we then have that  $F_\varepsilon(u_\varepsilon^*) \rightarrow F(u)$  as  $\varepsilon \rightarrow 0^+$ . This proves condition 2 in the  $\mathcal{C}^1$  case. To prove the general case we use a density argument. In Theorem 1.24 of [20] it is proved that every set with bounded perimeter can be approximated in  $L^1$  and in perimeter by a sequence of sets with  $\mathcal{C}^\infty$  boundaries. Thus, given any  $u \in BV(\mathcal{T}; \{\pm m_\beta\})$ , by a diagonalization procedure we then construct a sequence  $\{u_\varepsilon\}$  that satisfies condition 2. The  $\Gamma$ -convergence of  $\{F_\varepsilon\}$  to  $F$  is thus proved.

## APPENDIX A

In what follows we denote by  $C_\gamma(r)$  the atom of  $\mathcal{Q}_\gamma$  that contains  $r$  ( $\mathcal{Q}_\gamma$  is defined at the beginning of Section 3).

**Lemma A.1.** There is a positive constant  $c_1$  such that for any spin configuration  $\sigma_\gamma$ ,  $r \in \mathbb{R}^d$  and  $R > 0$  we have

$$\left| \int_{B_R(r)} dr' \sigma_\gamma(r') - \int_{B_R(r)} dr' \pi_\gamma \sigma_\gamma(r') \right| \leq c_1 R^{-1} \gamma^\delta$$

*Proof.* By definition, see (3.4),

$$\int_{B_R(r)} dr' J_\gamma^{(2)} = \int_{B_R(r)} dr' \frac{1}{|C_\gamma(r')|} \int_{C_\gamma(r')} dr'' \sigma_\gamma(r'')$$

Then

$$\begin{aligned} & \int_{B_R(r)} dr' \pi_\gamma \sigma_\gamma(r') - \int_{B_R(r)} dr' \sigma_\gamma(r') \\ &= \int_{B_R(r)} dr' \frac{1}{|C_\gamma(r')|} \int_{C_\gamma(r') \setminus (C_\gamma(r') \cap B_R(r))} dr'' \sigma_\gamma(r'') \\ & \quad - \int_{B_R(r)} dr' \frac{1}{|C_\gamma(r') \cap B_R(r)|} \\ & \quad \times \int_{C_\gamma(r') \cap B_R(r)} dr'' \sigma_\gamma(r'') \left[ 1 - \frac{|C_\gamma(r') \cap B_R(r)|}{|C_\gamma(r')|} \right] =: \text{I} + \text{II} \end{aligned}$$

Since  $C_\gamma(r') \cap B_R(r) = C_\gamma(r')$  for  $|r' - r| < R - 2\gamma^\delta$ , [recall (3.3)] there is a positive constant  $c_1$  such that  $\max(\text{I}, \text{II}) \leq c_1 R^{-1} \gamma^\delta$ . ■

**Lemma A.2.** There is a positive constant  $c_2$  such that for any spin configuration  $\sigma_\gamma$

$$|H_\varepsilon(\sigma_\gamma) - H_\varepsilon(\pi_\gamma \sigma_\gamma)| \leq c_2 \gamma^\delta \varepsilon^{-d} \quad (\text{A.1})$$

*Proof.* Recalling definition (2.3),  $H_\varepsilon(\cdot) = H(\cdot; \mathcal{J}_\varepsilon)$  and (3.4), we have

$$\begin{aligned}
& H_\varepsilon(\pi_\gamma \sigma_\gamma) - H_\varepsilon(\sigma_\gamma) \\
&= \frac{1}{2} \int_{\mathcal{J}_\varepsilon} dr_1 \int_{\mathcal{J}_\varepsilon} dr_2 J(|r_1 - r_2|) \sigma_\gamma(r_1) \sigma_\gamma(r_2) \\
&\quad - \frac{\gamma^{-2\delta d}}{2} \int_{\mathcal{J}_\varepsilon} dr \int_{\mathcal{J}_\varepsilon} dr' J(|r - r'|) \int_{C_\gamma(r)} dr_1 \int_{C_\gamma(r')} dr_2 \sigma_\gamma(r_1) \sigma_\gamma(r_2) \\
&= -\frac{1}{2} \int_{\mathcal{J}_\varepsilon} dr_1 \int_{\mathcal{J}_\varepsilon} dr_2 \sigma_\gamma(r_1) \sigma_\gamma(r_2) \left[ \gamma^{-2\delta d} \int_{C_\gamma(r_1)} dr \right. \\
&\quad \left. \times \int_{C_\gamma(r_2)} dr' J(|r - r'|) - J(|r_1 - r_2|) \right]
\end{aligned}$$

By the regularity of  $J$ , there is a positive constant  $c'$  so that

$$\sup_{\substack{r \in C_\gamma(r_1) \\ r' \in C_\gamma(r_2)}} |J(|r - r'|) - J(|r_1 - r_2|)| \leq c' \mathbf{1}_{\{|r_1 - r_2| \leq 2\}} \|\nabla J\|_\infty \gamma^\delta$$

where  $\|\nabla J\|_\infty$  is the sup norm of  $\nabla J$  and  $\mathbf{1}_A$  the characteristic function of  $A$ . We then get

$$|H_\varepsilon(\sigma_\gamma) - H_\varepsilon(\pi_\gamma \sigma_\gamma)| \leq c_2 \gamma^\delta \varepsilon^{-d} \quad \blacksquare$$

**Lemma A.3.** There is a positive constant  $c_3$  such that for any coarse grained configuration  $s_\gamma$

$$|\log W(s_\gamma) - \gamma^{-d} \mathcal{J}_\varepsilon(s_\gamma)| \leq c_3 (\gamma \varepsilon)^{-d} \gamma^{2(1-\delta)d} \log(\gamma^{-1})$$

*Proof.* Let  $C_\gamma$  be an atom of  $\mathcal{Q}_\gamma$  and  $r \in C_\gamma$ . Then, by definition,

$$s_\gamma(r) = \frac{1}{|C_\gamma(r)|} \int_{C_\gamma(r)} dr' \sigma_\gamma(r')$$

We call  $\lambda_1 \dots \lambda_N$  the values attained by  $\sigma_\gamma(r')$  when  $r'$  varies in  $C_\gamma(r)$ . Thus  $\lambda_i \in \{\pm 1\}$  and  $N$  goes like  $\gamma^{(\delta-1)d}$ . The number of sequences  $\{\lambda_i\}$  that give rise to the same value  $m \in [-1, 1]$  of  $s_\gamma(r)$  is

$$\binom{N}{K(m)}, \quad \text{where } K(m) := \frac{1-m}{2} N \in \mathbb{N}$$

We use the Stirling's formula to estimate this quantity, see for instance [3] and references therein, and we get, recalling the definition (2.21) of  $i(m)$ ,

$$\begin{aligned} \left| \log \binom{N}{K(m)} - i(m) \right| &\leq \frac{\log N}{2N} - \frac{1}{N} \log \frac{(1-m)(1+m)}{4} \\ &\quad + \frac{4}{N^2(1-m)(1+m)} \\ &\leq c'_3 \frac{\log N}{N} \end{aligned} \tag{A.2}$$

for  $m = 0, \pm 2/N, \dots, \pm(N-2)/N$  and for a suitable positive constant  $c'_2$ . The left hand side in (A.2) is 0 if  $m = \pm 1$ .

The weight  $W(s_\gamma)$  of the configuration  $s_\gamma$  is the product of the weights over all the cubes  $C_\gamma$  in  $\mathcal{T}_\varepsilon$ . We call  $C_i$  the generic one and  $m_i$  the value of  $s_\gamma(r)$  when  $r \in C_i$ . We then have

$$\left| \sum_{C_i} \log \binom{N}{K(m_i)} - \gamma^{-d} \int_{\mathcal{T}_\varepsilon} dr i(m(r)) \right| \leq c_3 (\gamma\varepsilon)^{-d} \gamma^{2d(1-\delta)} \log(\gamma^{-1})$$

because the number of terms in the sum is of order  $\varepsilon^{-d} \gamma^{-\delta d}$ . ■

**Lemma A.4.** Let  $N_\gamma$  be the total number of distinct coarse grained configurations  $s_\gamma$ ; then there is a positive constant  $c_4$  so that

$$\log N_\gamma \leq c_4 (\gamma\varepsilon)^{-d} \gamma^{(1-\delta)d} \log(\gamma^{-1})$$

*Proof.* A configuration  $s_\gamma$  is defined by giving the value of the magnetization in each cube  $C_\gamma$  paving  $\mathcal{T}_\varepsilon$ . Since the number of possible values of the magnetization is bounded by  $2\gamma^{(\delta-1)d}$  and the number of cubes  $C_\gamma$  in  $\mathcal{T}_\varepsilon$  is bounded by  $\varepsilon^{-d} \gamma^{-\delta d}$ , we immediately conclude the proof of the Lemma. ■

## APPENDIX B

In this Appendix we prove some statements used in the proof of Proposition 4.2. It is convenient to formulate the problem in the following way. We consider three non-empty, disjoint, bounded regions of  $\mathbb{R}^d$ ,  $\Delta$ ,  $\Gamma$  and  $\Lambda$ . Each one is union of cubes of the partition  $\mathcal{Q}^{(-1)}$ . We suppose that  $\mathcal{R} := \Delta \cup \Gamma \cup \Lambda$  is connected and that  $d(\Delta, \Lambda)$ , the Euclidean distance between the two sets, is not smaller than 2.




Fig.4. An example of the regions  $\Delta$ ,  $\Gamma$  and  $\Lambda$  of Appendix B.  $\Delta$  is the union of the three central cubes,  $\Gamma$  of the ten cubes surrounding  $\Delta$  and  $\Lambda$  is the complement. The shadowed region represents  $\Delta_\ell$ .

The example to have in mind is with  $\Delta$  a connected set,  $\Gamma$  the collection of all the cubes in  $\mathcal{R}$  outside  $\Delta$  that have distance 0 from  $\Delta$ ;  $\Lambda$  is then separated by  $\Delta$  at least by a cube of  $\mathcal{Q}^{(-1)}$ , whose side has length 2 (see Figure 4).

We consider the functional  $\mathcal{F}(m; \mathcal{R})$  on  $L^\infty(\mathcal{R}; [-m_\beta, m_\beta])$  defined in the usual way by

$$\begin{aligned} \mathcal{F}(m; \mathcal{R}) := & \int_{\mathcal{R}} dr [f(m(r)) - f(m_\beta)] \\ & + \frac{1}{4} \iint_{\mathcal{R} \times \mathcal{R}} dr dr' J(|r - r'|) [m(r) - m(r')]^2 \end{aligned} \quad (\text{B.1})$$

Given  $\ell \in (0, 1)$ , we set

$$\Delta_\ell := \{r \in \Delta : d(r, \Gamma) \leq \ell\}; \quad \Gamma_\ell := \{r \in \Gamma : d(r, \Delta) \leq 1 - \ell\} \quad (\text{B.2})$$

and finally, for any  $r \in \mathcal{R}$ , any measurable set  $C \subset \mathcal{R}$  and  $m \in L^\infty(\mathcal{R}; [-m_\beta, m_\beta])$ , we define

$$h(r; m; C) := \frac{\int_C dr' J(|r - r'|) m(r')}{\int_C dr' J(|r - r'|)} \quad (\text{B.3})$$

setting  $h(r; m; C) := 1$  if the denominator in (B.3) is 0.

We will consider two cases:  $C = \Gamma \cup \Lambda$  and  $r \in \Delta_\ell \cup \Gamma_\ell$  and  $C = (\Gamma \setminus \Gamma_\ell) \cup \Lambda$  and  $r \in \Gamma_\ell$ . In both cases  $h(r; m; C)$  does not depend on the values of  $m$  in  $\Lambda$  and, moreover, the denominator in (B.3) is uniformly positive because, by Definition 2.1c,  $\sup\{s > 0 : J(s) > 0\} = 1$ . This remark is used to prove the following lemma:

**Lemma B.1.** For any  $\ell \in (0, 1)$  there exists a sequence  $\{c_k\}_{k \in \mathbb{N}}$  of positive numbers which converges to 0 as  $k \rightarrow +\infty$  with the following property. If  $\zeta > 0$  and for some  $k \in \mathbb{N}$

$$\pi^{(k)}m(r) \geq m_\beta - \zeta \quad \text{for all } r \in \Gamma \quad (\text{B.4})$$

then

$$h(r; m; \Gamma \cup \Lambda) \geq m_\beta - (\zeta + c_k) \quad \text{for all } r \in \Delta_\ell \cup \Gamma_\ell \quad (\text{B.5})$$

$$h(r; m; (\Gamma \setminus \Gamma_\ell) \cup \Lambda) \geq m_\beta - (\zeta + c_k) \quad \text{for all } r \in \Gamma_\ell \quad (\text{B.6})$$

*Proof.* Let  $C = \Gamma \cup \Lambda$  and

$$c_k := \sup_{r \in \Delta_\ell \cup \Gamma_\ell} \sup_{m \in L^\infty(\mathcal{R}; [-m_\beta, m_\beta])} |h(r; m; C) - h(r; \pi^{(k)}m; C)| \quad (\text{B.7})$$

Then  $c_k \rightarrow 0$  as  $k \rightarrow +\infty$  and if  $\pi^{(k)}m(r) \geq m_\beta - \zeta$  for  $r \in \Gamma$  then

$$h(r; m; C) \geq m_\beta - \zeta - c_k \quad \text{for all } r \in \Delta_\ell \cup \Gamma_\ell$$

that proves (B.5). The proof of (B.6) is similar.

**Remark.** We will actually take for  $c_k$  the maximum over all possible choices of  $\Delta$  and  $\Gamma$  (which is finite because the right hand side of (B.7) takes only finitely many values).

For any  $m \in L^\infty(\mathcal{R}; [-m_\beta, m_\beta])$ ,  $\ell \in (0, 1)$  and  $\theta \in (0, m_\beta)$  we define

$$m_{\ell, \theta}(r) := \begin{cases} m(r) & \text{if } r \in (\Gamma \setminus \Gamma_\ell) \cup \Lambda \\ |m(r)| & \text{if } r \in \Delta \cup \Gamma_\ell \text{ and } |m(r)| \geq m_\beta - \theta \\ m_\beta - \theta & \text{if } r \in \Delta \cup \Gamma_\ell \text{ and } |m(r)| < m_\beta - \theta \end{cases} \quad (\text{B.8})$$

We then have

**Theorem B.2.** Let  $\ell \in (0, 1)$ ,  $\zeta > 0$  and  $k \in \mathbb{N}$  be such that  $\zeta + c_k =: \theta < m_\beta$ , with  $c_k$  defined in (B.7). Then for all  $m$  that satisfy (B.4) we have

$$\mathcal{F}(m; \mathcal{R}) \geq \mathcal{F}(m_{\ell, \theta}; \mathcal{R}) \quad (\text{B.9})$$

*Proof.* Since  $m_{\ell, \theta} = m$  in  $(\Gamma \setminus \Gamma_\ell) \cup \Lambda$  and the support of  $J$  is contained in the unit ball, there is no interaction between  $\Lambda$  and  $\Delta \cup \Gamma_\ell$ , where  $m$  and

$m_{\ell,\theta}$  are different. For this reason the values of  $m$  in  $\Lambda$  will not play any role in the following proof.

We observe that under the replacement  $m \rightarrow m_{\ell,\theta}$  the first term in the expression of  $\mathcal{F}(\cdot; \mathcal{R})$  does not increase, so it is sufficient to prove that the second term has the same property. Given two bounded measurable sets  $A$  and  $B$  in  $\mathbb{R}^d$  we write

$$E(A, B) = \frac{1}{4} \int_{r \in A} \int_{r' \in B} dr dr' J(|r - r'|) \times \{ [m_{\ell,\theta}(r) - m_{\ell,\theta}(r')]^2 - [m(r) - m(r')]^2 \} \quad (\text{B.10})$$

and split the second term in (B.1) into a sum of terms  $E(A_i, B_i)$ . We will show that each of them is non-positive and this will prove the Theorem. In the proof we will use the two following statements whose simple proof is omitted:

For any  $s \in [-m_\beta, m_\beta]$  and  $C \subset \mathcal{R}$  define

$$G(s, C) := \int_C dr' J(|r - r'|) [s - m(r')]^2$$

Then

1.  $G(s, C) \geq G(t, C)$  for any  $s \leq t \leq h(r; m; C)$ .
2. If  $h(r; m; C) > 0$  then  $G(s, C) \geq G(|s|, C)$  for any  $s \in [-m_\beta, m_\beta]$ .

We next examine separately the various terms  $E(A_i, B_i)$ :

(T1).  $A = B = (\Gamma \setminus \Gamma_\ell) \cup \Lambda$ . Then  $E(A, B) = 0$  because  $m = m_{\ell,\theta}$ .

(T2).  $A = \Gamma_\ell \cap \{|m(r)| < m_\beta - \theta\}$  and  $B = (\Gamma \setminus \Gamma_\ell) \cup \Lambda$ . We apply statement 1 for any  $r \in A$  with  $s := m(r)$  and  $t := m_{\ell,\theta}(r) = m_\beta - \theta$ . In fact  $s \leq t \leq h(r; m; B)$ , the last inequality follows from (B.6) recalling that  $\theta = \zeta + c_k$ , see (B.6). We then have  $G(m(r), B) \geq G(m_{\ell,\theta}(r), B)$ , which, integrated over  $r \in A$ , yields  $E(A, B) \leq 0$ .

(T3).  $A = \Gamma_\ell \cap \{|m(r)| \geq m_\beta - \theta\}$  and  $B = (\Gamma \setminus \Gamma_\ell) \cup \Lambda$ . For any  $r \in A$  we have  $h(r; m; B) \geq m_\beta - \theta > 0$ . Since  $m_{\ell,\theta} = |m|$  on  $A$ , we get by statement 2 that  $G(m(r), B) \geq G(m_{\ell,\theta}(r), B)$ . Integrating this inequality over  $r \in A$  we then find that  $E(A, B) \leq 0$ .

(T4).  $A = \Delta_\ell \cap \{|m(r)| < m_\beta - \theta\}$  and  $B = \Gamma \cup \Lambda$ . We write

$$\begin{aligned} E(A, B) &= \frac{1}{4} \int_A dr \int_B dr' J(|r - r'|) \\ &\quad \times \{ ([m_{\ell,\theta}(r) - m(r')]^2 - [m(r) - m(r')]^2) \\ &\quad + ([m_{\ell,\theta}(r) - m_{\ell,\theta}(r')]^2 - [m_{\ell,\theta}(r) - m(r')]^2) \} \\ &=: \text{I}(A, B) + \text{II}(A, B) \end{aligned} \quad (\text{B.11})$$

For any  $r \in A$  we have  $h(r; m; B) \geq m_\beta - \theta$  by (B.5) and  $m_\beta - \theta = m_{\ell, \theta}(r) =: t \geq s := m(r)$ . Hence reasoning as in (T2) we get  $\text{I}(A, B) \leq 0$ .

To prove that  $\text{II}(A, B) \leq 0$  we introduce the sets:

$$\begin{aligned} B_1 &:= \{r' \in \Gamma_\ell : |m(r')| < m_\beta - \theta\} \\ B_2 &:= \{r' \in \Gamma_\ell : |m(r')| \geq m_\beta - \theta\} \\ B_3 &:= \Gamma \setminus \Gamma_\ell \end{aligned}$$

and we split  $\text{II}(A, B) = \text{II}(A, B_1) + \text{II}(A, B_2) + \text{II}(A, B_3)$ . Then  $m_{\ell, \theta}(r') = m(r')$  for any  $r' \in B_3$  so that  $\text{II}(A, B_3) = 0$ . Moreover  $m_{\ell, \theta}(r') = m_\beta - \theta$  for  $r' \in B_1$  and  $m_{\ell, \theta}(r) \geq m_\beta - \theta$  for  $r \in \Delta_\ell$ , so that  $|m_{\ell, \theta}(r) - m_{\ell, \theta}(r')| \leq |m_{\ell, \theta}(r) - m(r')|$  for  $r \in A$  and  $r' \in B_1$ . Therefore  $\text{II}(A, B_1) \leq 0$ . Finally we have  $m_{\ell, \theta}(r') = |m(r')|$  for  $r' \in B_2$ . Hence

$$\begin{aligned} \text{II}(A, B_2) &= \frac{1}{4} \int_{r' \in B_2} dr' \int_{r \in A} dr J(|r - r'|) \\ &\quad \times \{[|m(r')| - m_{\ell, \theta}(r)]^2 - [m(r') - m_{\ell, \theta}(r)]^2\} \leq 0 \end{aligned}$$

by statement 2 because  $h(r'; m_{\ell, \theta}; A) \geq m_\beta - \theta > 0$  for any  $r' \in B_2$  as  $m_{\ell, \theta} \geq m_\beta - \theta$ .

(T5).  $A = \Delta_\ell \cap \{|m(r)| \geq m_\beta - \theta\}$  and  $B = \Gamma \cup \Lambda$ . We use (B.11) with the new  $A$ . The first term on the right hand side is non-negative by statement 2: in fact  $h(r; m; B) > 0$  for  $r \in A$  by (B.5) and  $m_{\ell, \theta} = |m|$  on  $A$  by (B.8). For the second term we use the same argument as in the last part of the proof of (T4). Thus  $E(A, B) \leq 0$ .

(T6).  $A = B = \Gamma_\ell$ ,  $A = B = \Delta$  and  $A = \Delta \setminus \Delta_\ell$ ,  $B = \Gamma_\ell$ . In all these cases  $|m_{\ell, \theta}(r) - m_{\ell, \theta}(r')| \leq |m(r) - m(r')|$  for any  $r \in A$ ,  $r' \in B$ . Hence  $E(A, B) \leq 0$ .

The Theorem is proved.  $\blacksquare$

## APPENDIX C

In this Appendix we prove the properties of the instanton used in the proof of Proposition 4.2. Recall that  $K_\varepsilon$  is a cylinder in  $\mathbb{R}^d$  with basis  $B_\varepsilon = \varepsilon^{-1}B$  and that we have supposed for simplicity that  $K_\varepsilon$  is  $\mathcal{Q}^{(0)}$ -measurable. The case when  $K_\varepsilon$  is a torus has been considered in [24]. We will first prove the existence of  $m_\varepsilon^{(7)}$ , see the proof of Proposition 4.2, hereafter denoted for simplicity by  $\tilde{m}$ , namely of an instanton solution of (4.19). We will follow



[14] where the  $d = 1$  case is considered. We use the basic and elementary properties of the evolution (4.19) listed below, where  $m_t$ ,  $t \geq 0$ , denotes a solution of (4.19) with  $m_0 \in L^\infty(K_\varepsilon; [-1, 1])$ .

1. Let  $\psi_t := m_t - e^{-t}m_0$ . Then  $|\nabla\psi_t|$  is uniformly bounded with respect to  $r$ ,  $t$  and  $m_0$ . As a consequence there exists a sequence  $t_n \rightarrow +\infty$  such that  $m_{t_n}$  converges uniformly on the compact subsets of  $K_\varepsilon$ , as  $n \rightarrow +\infty$ .
2. The functional  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon)$  defined in (4.16) is lower semicontinuous on  $L^\infty(K_\varepsilon; [-1, 1])$  in the weak  $L^2_{\text{loc}}(K_\varepsilon)$ -topology, and  $\mathcal{F}^{(1)}(m; K_\varepsilon) < +\infty$  if and only if there are  $\sigma_\pm \in \{\pm 1\}$ , such that  $m - \chi_\sigma \in L^2(K_\varepsilon, dr)$ , where  $\chi_\sigma(r) := m_\beta \sigma_+$  (respectively  $\chi_\sigma(r) := m_\beta \sigma_-$ ) if  $r_d \geq 0$  (respectively  $r_d < 0$ ).
3. If  $\mathcal{F}^{(1)}(m_0; K_\varepsilon) < +\infty$ , then  $d\mathcal{F}^{(1)}(m_t; K_\varepsilon)/dt \leq 0$  for all  $t \geq 0$ .
4. If  $\mathcal{F}^{(1)}(m_0; K_\varepsilon) < +\infty$ , then there are  $\hat{m} \in \mathcal{C}(K_\varepsilon; [-m_\beta, m_\beta])$  and a sequence  $t_n \rightarrow +\infty$  such that  $m_{t_n} \rightarrow \hat{m}$  uniformly on the compact subsets of  $K_\varepsilon$  as  $n \rightarrow +\infty$ , and  $\hat{m}$  is a stationary solution of (4.19) in the whole  $K_\varepsilon$ .

We omit the proof of the above properties which is very similar to that in [14] for  $d = 1$ .

**Proposition C.1.** There is  $\tilde{m} \in \mathcal{C}^\infty(K_\varepsilon; [-m_\beta, m_\beta])$  which is a stationary solution of (4.19). Moreover  $\tilde{m}$  is an antisymmetric and strictly increasing function of  $r_d$ .

*Proof.* Following [14], we set for  $r \in K_\varepsilon$

$$m_0(r) := \begin{cases} -m_\beta & \text{for } r_d \leq -1 \\ m_\beta & \text{for } r_d \geq 1 \\ m_\beta r_d & \text{otherwise} \end{cases} \quad (\text{C.1})$$

Let  $m_t$  solve (4.19) with initial datum (C.1). Then  $m_t$  is non-decreasing and antisymmetric for any  $t \geq 0$ . Since  $\mathcal{F}^{(1)}(m_0; K_\varepsilon) < +\infty$ , the limit  $\hat{m}$  considered in property 4 solves (C.2) below, is non-decreasing, antisymmetric and such that  $\mathcal{F}^{(1)}(\hat{m}; K_\varepsilon) < +\infty$ . This shows that  $\tilde{m} := \hat{m}$  is not identically 0.

Recalling that  $J^{(1)}$ , defined in (4.20), is a  $\mathcal{C}^\infty$  function, we then obtain by differentiating the equation

$$\tilde{m}(r) = \tanh\{\beta J^{(1)} * \tilde{m}(r)\}, \quad r \in K_\varepsilon \quad (\text{C.2})$$

that  $\tilde{m} \in \mathcal{C}^\infty(K; [-m_\beta, m_\beta])$ . If the derivative  $\tilde{m}'$  of  $\tilde{m}(r)$  with respect to  $r_d$  were 0 at some point  $r$ , then it would be 0 in the whole  $\{r' : J^{(1)}(r, r') > 0\}$  because  $\tilde{m}' \geq 0$ . By iteration we would then reach a contradiction with the previous statement that  $\tilde{m}$  is not identically 0. Hence  $\tilde{m}(r)$  is a strictly increasing function of  $r_d$  and the Proposition is proved.  $\blacksquare$

**Proposition C.2.** There are constants  $c$  and  $c'$  positive and independent of  $\varepsilon$  so that the function  $\tilde{m}$  verifies the inequality

$$\tilde{m}(r) \geq m_\beta - c'e^{-cr_d}, \quad r_d \geq 0 \quad (\text{C.3})$$

*Proof.* By Proposition C.1 there is  $a > 0$  so that  $\tilde{m} \geq a$  on  $\{r \in K_\varepsilon : r_d \geq 1\}$ . Using this fact we will prove that (C.3) holds with a constant  $c'$  that depends on  $a$ . We will then complete the proof of the Proposition by showing that  $a$  (hence  $c'$ ) is uniformly bounded away from zero as  $\varepsilon \rightarrow 0^+$ .

To prove the first statement we introduce the following dynamics. We call  $K_\varepsilon^{(i)} := \{r \in K_\varepsilon : r_d \geq i\}$ ,  $i = 1, 2$ . Then for any  $m \in L^\infty(K_\varepsilon^{(1)}; [-1, 1])$  we define  $m_t \in L^\infty(K_\varepsilon^{(1)}; [-1, 1])$  as  $m_0 := m$  on  $K_\varepsilon^{(1)}$ ,  $m_t = m_0$ ,  $t \geq 0$ , on  $K_\varepsilon^{(1)} \setminus K_\varepsilon^{(2)}$ . Finally, on  $K_\varepsilon^{(2)}$ , for  $t \geq 0$   $m_t$  solves the equation

$$\begin{aligned} \frac{dm_t(r)}{dt} &= -m_t(r) + \tanh\{\beta J^{(1)} * m_t(r)\} \\ J^{(1)} * m_t(r) &= \int_{K_\varepsilon^{(1)}} dr' J^{(1)}(r, r') m_t(r') \end{aligned} \quad (\text{C.4})$$

$\tilde{m}$  restricted to  $K_\varepsilon^{(1)}$  is obviously a stationary solution of (C.4).

Given  $u_0 \in L^\infty(K_\varepsilon^{(1)}; [-1, 1])$  we say that the function  $u_t \in L^\infty(K_\varepsilon^{(1)}; [-1, 1])$  is a subsolution of (C.4) with initial datum  $u_0$  if  $u_t = u_0$ ,  $t \geq 0$ , on  $K_\varepsilon^{(1)} \setminus K_\varepsilon^{(2)}$  and

$$\frac{du_t}{dt} \leq -u_t + \tanh\{\beta J^{(1)} * u_t\} \quad \text{on } K_\varepsilon^{(2)} \text{ for all } t \geq 0 \quad (\text{C.5})$$

Since  $J^{(1)} \geq 0$  one can show that if  $u_0 \leq m_0$  on  $K_\varepsilon^{(1)}$ , then  $u_t \leq m_t$  on  $K_\varepsilon^{(1)}$  for all  $t \geq 0$ . A similar statement is proved in [14] for  $d = 1$ .

We want to construct a subsolution of (C.4). We start by constructing a countable system of functions  $v_t : \mathbb{N}_+ \rightarrow \mathbb{R}$ ,  $t \geq 0$ , defined by  $v_t(1) := a$ ,  $t \geq 0$ , and, for  $n \in \mathbb{N}$ ,  $n \geq 1$ ,

$$\frac{dv_t(n+1)}{dt} = -v_t(n+1) + \tanh\{\beta v_t(n)\}, \quad v_0(n+1) = a \quad (\text{C.6})$$

i.e.

$$v_t(n+1) = e^{-t}a + \int_0^t ds e^{-(t-s)} \tanh\{\beta v_s(n)\} \quad (\text{C.7})$$

Since  $\beta > 1$ , we have  $a - \tanh(\beta a) < 0$  (recall that  $a < m_\beta$ ) so that

$$v_t(2) = \tanh(\beta a) + e^{-t}[a - \tanh(\beta a)] \quad (\text{C.8})$$

is an increasing function of  $t$ . Define

$$v^*(2) := \lim_{t \rightarrow \infty} v_t(2) = \tanh\{\beta a\} \quad (\text{C.9})$$

Similarly, by induction on  $n$ , setting

$$v^*(n+1) := \lim_{t \rightarrow \infty} v_t(n+1)$$

one can prove that

$$v^*(n+1) = \tanh\{\beta v^*(n)\} \quad (\text{C.10})$$

Calling  $T(s) := \tanh\{\beta s\}$  the map from  $(0, m_\beta]$  into itself and  $T^n$  the  $n$ -th iterate of  $T$ , we have from (C.10):

$$v^*(n+1) = T^n(a)$$

Hence there are constants  $c_1 > 0$  and  $c_2$  so that

$$v^*(n) \geq m_\beta - c_2 e^{-c_1 n} \quad (\text{C.11})$$

because  $m_\beta$  is a stable point for  $T$  attracting any orbit that starts from  $(0, m_\beta]$ .

To relate the system  $v_t(n)$ ,  $n \in \mathbb{N}$ , to a subsolution of (C.4) we need preliminarily to prove that  $D_t(n+1) := v_t(n+1) - v_t(n) \geq 0$  for all  $t \geq 0$ . We have, after Taylor expanding to first order,

$$\begin{aligned} D_t(n+1) &= \int_0^t ds e^{-(t-s)} (\tanh\{\beta v_s(n)\} - \tanh\{\beta v_s(n-1)\}) \\ &= \int_0^t ds e^{-(t-s)} \psi_s(n) D_s(n) \end{aligned} \quad (\text{C.12})$$

where  $\psi_s(n)$  is the derivative of the hyperbolic tangent computed at some value which depends on  $v_s(n-1)$  and  $v_s(n)$ . Then  $\psi_s(n) > 0$  and, consequently,  $D_t(n+1) \geq 0$  for all  $n \geq 1$  and all  $t \geq 0$ , because  $D_t(2) \geq 0$  as it follows from (C.8) recalling that  $v_t(1) = a$ .

We are going to show that the function  $u_t \in L^\infty(K_\varepsilon^{(1)}; [a, m_\beta])$ ,  $t \geq 0$ , defined by

$$u_t(r) := v_t([r_d]) \quad (\text{C.13})$$

is a subsolution of (C.4). In fact let  $r \in K_\varepsilon^{(2)}$  be such that  $[r_d] = n + 1$ ; then

$$\begin{aligned} \frac{du_t(r)}{dt} + u_t(r) - \tanh\{\beta J^{(1)} * u_t(r)\} \\ = -\tanh\{\beta J^{(1)} * u_t(r)\} + \tanh\{\beta u_t(n)\} \leq 0 \end{aligned} \quad (\text{C.14})$$

because for such values of  $r$  one has

$$J^{(1)} * u_t(r) = k_1(r)v_t(n+2) + k_2(r)v_t(n+1) + k_3(r)v_t(n) \geq v_t(n) \quad (\text{C.15})$$

for three suitable non-negative functions  $k_1$ ,  $k_2$  and  $k_3$ . The first equality follows from the fact that  $J^{(1)}(r, r') = 0$  if  $|r - r'| \geq 1$ . Moreover, recalling that the  $r'$ -integral of  $J^{(1)}$  is equal to 1, we get  $k_1(r) + k_2(r) + k_3(r) \equiv 1$ . The last inequality in (C.15) follows from the fact that  $v_t(m) \geq v_t(n)$  for  $m \geq n$ , which has already been proved. We have thus shown that  $u_t$  satisfies (C.5) and that it is a subsolution of (C.4). Since  $\tilde{m} \geq u_0$  on  $K_\varepsilon^{(1)}$ , it then follows that

$$\tilde{m} \geq \lim_{t \rightarrow +\infty} u_t = u^*, \quad u^*(r) := v^*([r_d])$$

(C.3) then follows from (C.11), but, as already observed, the constant  $c'$  depends on  $c_2$  which is in turn determined by  $a$ . To prove Proposition C.2 we thus need to show that  $a$  can be taken independent of  $\varepsilon$ .

To this end for any  $r \in K_\varepsilon$  with  $1 \leq r_d \leq 2$  we consider a region  $\Gamma \subset K_\varepsilon \cap \{1 \leq r_d \leq 2\}$  containing  $r$  which is  $\mathcal{Q}^{(0)}$ -measurable and  $*$ -connected, see Definition 2.1a and Definition 4.3. We also suppose that for  $\alpha > 0$  as in (C.21) below,

$$\frac{|\delta\Gamma|}{|\Gamma|} \leq \alpha \quad (\text{C.16})$$

$|\Gamma|$  is the volume of  $\Gamma$ , hence the number of cubes of  $\mathcal{Q}^{(0)}$  in  $\Gamma$ ;  $\delta\Gamma$  is the union of all the cubes in  $K_\varepsilon \setminus \Gamma$  at distance not larger than 1 from  $\Gamma$  and  $|\delta\Gamma|$  the total number of such cubes. We can find a finite family of such sets  $\Gamma$  so that, modulo translations, for any  $\varepsilon > 0$  any  $r$  in the strip  $\{1 < r_d \leq 2\} \cap K_\varepsilon$  is contained in an element of the family. We fix in the sequel a region  $\Gamma$  and call  $\Lambda = \Gamma \cup \delta\Gamma$ .

Let  $u \in L^\infty(\Lambda; \{0, m_\beta\})$ ,  $u(r) := m_\beta$  when  $r \in \Gamma$  and  $u_0(r) := 0$  when  $r \in \delta\Gamma$ . We call  $u_t \in L^\infty(\Lambda; [0, m_\beta])$ ,  $t \geq 0$ , the function that solves (C.4) in  $\Gamma$ , with  $u_0 = u$  and  $u_t = 0$ , on  $\delta\Gamma$  for all  $t \geq 0$ .

The Proposition is then a consequence of the following:

**Lemma.** (i) There are  $u^* \in L^\infty(\Lambda; [0, m_\beta])$  verifying (C.2) in  $\Gamma$  and a sequence  $t_n \rightarrow +\infty$  as  $n \rightarrow +\infty$  such that  $u_{t_n} \rightarrow u^*$  uniformly on  $\Lambda$ .

(ii)  $\tilde{m} \geq u^*$  on  $\Lambda$ .

(iii) There is  $a > 0$  so that  $u^* \geq a$  on  $\Gamma$ .

*Proof of the Lemma.* Statement (i) follows from property 4, see the beginning of this Appendix, that also applies to  $u_t(r)$ .

Let  $m_t(r)$  be the solution of (4.19) with initial datum (C.1). Then  $m_t \geq 0$  on  $\delta\Gamma$  because  $m_t \geq 0$  on  $\{r_d \geq 0\}$ , as  $m_t$  is antisymmetric and non-decreasing. It then follows that  $m_t \geq u_t$ ,  $t \geq 0$  on  $\Lambda$ .

The proof of (iii) is more delicate. We know from (i) that for  $r \in \Gamma$

$$u^*(r) = \tanh \left\{ \int_{\Lambda} dr' J^{(1)}(r, r') u^*(r') \right\} \quad (\text{C.17})$$

Then there is a constant  $c_5 > 0$  such that, for any  $r \in \Gamma$ ,

$$u^*(r) \geq c_5 \int_{\Gamma} dr' J^{(1)}(r, r') u^*(r') \quad (\text{C.18})$$

Moreover there are a positive integer  $n$  and  $a' > 0$  so that for all  $r$  and  $r'$  in  $\Gamma$

$$\int_{\Gamma} dr_1 J^{(1)}(r, r_1) \cdots \int_{\Gamma} dr_{n-1} J^{(1)}(r_{n-2}, r_{n-1}) J^{(1)}(r_{n-1}, r') \geq a'$$

thus

$$u^*(r) \geq (c_5)^n a' \int_{\Gamma} dr' u^*(r'), \quad \text{for all } r \in \Gamma$$

It is therefore sufficient to prove that  $u^*$  is not identically 0. To this end we use a Liapunov function for the evolution satisfied by  $u_t$ . We set for  $u \in L^\infty(\Lambda; [0, m_\beta])$

$$\begin{aligned} \mathcal{F}^{(3)}(u; \Lambda) &:= \int_{\Gamma} dr [1 - j(r)] [f(u(r)) - f(m_\beta)] \\ &\quad + \frac{1}{4} \int_{\Gamma} dr \int_{\Gamma} dr' J(|r - r'|) [u(r) - u(r')]^2 \\ &\quad + \frac{1}{2} \int_{\Gamma} dr \int_{\delta\Gamma} dr' J(|r - r'|) [u(r) - u(r')]^2 \end{aligned}$$

One can check that  $\mathcal{F}^{(3)}(u_t; \Gamma)$  is a non-increasing function of  $t$  and that

$$\mathcal{F}^{(3)}(u^*; \Lambda) \leq \mathcal{F}^{(3)}(u_0; \Lambda) \quad (\text{C.19})$$

We will conclude that  $u^* \neq 0$ , 0 is here the function identically zero, from the fact that

$$\mathcal{F}^{(3)}(0; \Lambda) > \mathcal{F}^{(3)}(u_0; \Lambda) \quad (\text{C.20})$$

To prove (C.20) we observe that there are positive constants  $c_3$  and  $c_4$  so that

$$\begin{aligned} \mathcal{F}^{(3)}(0; \Lambda) &= \int_{\Gamma} dr [1 - j(r)] [f(0) - f(m_\beta)] \geq c_3 |\Gamma| \\ \mathcal{F}^{(3)}(u_0; \Gamma) &\leq \frac{1}{2} \int_{\Gamma} dr \int_{\Lambda \setminus \Gamma} dr' J(|r - r'|) m_\beta^2 \leq c_4 |\delta\Gamma| \end{aligned}$$

Then (C.20) follows from the inequality  $c_3 |\Gamma| > c_4 |\delta\Gamma|$ , which holds if we choose

$$\alpha \equiv \frac{|\delta\Gamma|}{|\Gamma|} = \frac{c_3}{2c_4} \quad (\text{C.21})$$

The Lemma and the Proposition are therefore proved.  $\blacksquare$

To prove that the instanton  $\tilde{m}$  is unique, modulo vertical translations and reflections, and to prove its stability, we follow again the approach used in [14]. We thus start from the linearization of (4.19) around  $\tilde{m}$ . Recalling (4.20),

$$\frac{dv}{dt} = Lv, \quad Lv(r) := -v(r) + \frac{1 - \tilde{m}(r)^2}{1 - j(r)} \beta J * v(r) \quad (\text{C.22})$$

$L$  is a self-adjoint operator in  $L^2(d\nu; K_\varepsilon)$  where

$$d\nu(r) = dr \frac{1 - j(r)}{1 - \tilde{m}(r)^2} \quad (\text{C.23})$$

Its spectrum contains the origin, as the  $r_d$ -derivative,  $\tilde{m}'$ , of  $\tilde{m}$  is a 0-eigenvector,

$$L\tilde{m}' = 0 \quad (\text{C.24})$$

We have proved in Proposition C.2 that  $\tilde{m}'$  is in  $L^2$  since it decays exponentially fast.

The spectrum of  $L$  is contained in the negative real axis. To prove it, we consider the Perron-Frobenius isomorphism  $U : L^2(d\nu; K_\varepsilon) \rightarrow L^2(d\mu; K_\varepsilon)$

$$Uv = \psi \quad \text{defined by} \quad U^{-1}\psi := \tilde{m}'\psi \quad (\text{C.25})$$

where

$$d\mu(r) := dr \tilde{m}'(r)^2 \frac{1 - j(r)}{1 - \tilde{m}(r)^2} \quad (\text{C.26})$$

The image of  $L$  under this isomorphism is

$$\mathcal{L} = ULU^{-1} \quad (\text{C.27})$$

which is the self-adjoint Markov generator

$$\begin{aligned} \mathcal{L}\psi(r) &= \int_{K_\varepsilon} dr' P(r, r') [\psi(r) - \psi(r')] \\ P(r, r') &:= [1 - \tilde{m}(r)^2] \frac{\tilde{m}'(r')}{\tilde{m}'(r)} \beta J^{(1)}(r, r') \end{aligned} \quad (\text{C.28})$$

The spectrum of  $\mathcal{L}$  lies therefore in the negative real axis, the same being true for its isomorphic image  $L$ . Denoting by  $\langle \cdot, \cdot \rangle_\mu$  the scalar product in  $L^2(d\mu; K_\varepsilon)$ , we have

$$\langle \psi, L\psi \rangle_\mu = -\frac{1}{2} \int_{K_\varepsilon} d\mu(r) \int_{K_\varepsilon} dr' P(r, r') [\psi(r) - \psi(r')]^2 \quad (\text{C.29})$$

which shows that 0 is a simple eigenvector, as (C.29) is equal to 0 only if  $\psi = c$ , a constant; then  $U^{-1}\psi = c\tilde{m}'$ .

We next show that there is a spectral gap, namely that 0 is an isolated eigenvalue and the rest of the spectrum is at finite distance from 0. We will use Weyl's theorem, [28], and to this end we decompose  $L = L_0 + L_1$  where

$$L_0 v := -v + \beta[1 - m_\beta^2] J^{(1)} * v \quad (\text{C.30})$$

$$L_1 v := \beta[\tilde{m}^2 - m_\beta^2] J^{(1)} * v \quad (\text{C.31})$$

We are going to prove that the spectrum of  $L_0$  is contained in  $\{\text{Re } z \leq -1 + \beta[1 - m_\beta^2]\}$ , recall that  $\beta[1 - m_\beta^2] < 1$ , because of the definition of  $m_\beta$ . It will suffice to prove that the spectral radius of  $1 + L_0$  is not larger than  $\beta[1 - m_\beta^2]$ . To see this we write

$$\begin{aligned} &\langle [1 + L_0]^n v, [1 + L_0]^n v \rangle_\mu \\ &\leq \frac{\|\tilde{m}'\|_\infty^2}{1 - m_\beta^2} \langle [1 + L_0]^n v, [1 + L_0]^n v \rangle_\lambda, \quad d\lambda(r) = dr[1 - j(r)] \end{aligned} \quad (\text{C.32})$$

and observe that  $L_0$  is self-adjoint in  $L^2(d\lambda; K)$  and

$$(1 + L_0)v(r) = \int_K dr' N(r, r')v(r'), \quad N(r, r') = \beta(1 - m_\beta^2)J^{(1)}(r, r') \quad (\text{C.33})$$

We then have, calling  $c := \|\tilde{m}'\|_\infty^2/(1 - m_\beta^2)$ ,

$$\begin{aligned} & \langle [1 + L_0]^n v, [1 + L_0]^n v \rangle_\lambda \\ & \leq c \int_K d\lambda(r) \int_K dr' N^{2n}(r, r') \frac{1}{2} [v(r)^2 + v(r')^2] \\ & \leq c [\beta(1 - m_\beta^2)]^{2n} \langle v, v \rangle_\lambda \end{aligned} \tag{C.34}$$

Thus  $\beta(1 - m_\beta^2)$  bounds the spectral radius of  $1 + L_0$  and this completes the proof of the statements concerning  $L_0$ . On the other hand  $L_1$  is a compact operator because it is an integral operator and because, by Proposition C.2,  $\tilde{m}(r)$  converges to  $m_\beta$  exponentially fast as  $|r_d| \rightarrow +\infty$ . By applying the Weyl's theorem we prove the spectral gap property for  $L$ , see [13] for more details.

We have therefore proved that the linearized evolution attracts towards the eigenvector  $\tilde{m}'$ . Proceeding as in [13] it is possible to prove the local stability of the manifold of instantons, namely the following property. Let  $m_0 \in L^\infty(K_\varepsilon; [-1, 1])$ . Call  $\tilde{m}_a$  the  $r_d$ -upward translation by  $a$  of  $\tilde{m}$ . Let  $\mu_a$  be the measure defined in (C.26) with  $\tilde{m}$  replaced by  $\tilde{m}_a$ . Suppose that there is  $a$  so that  $m_0 - \tilde{m}_a$  is in  $L^2(\mu_a; K_\varepsilon)$  and that its norm is smaller than some suitably fixed value. Then there exists  $b$  so that  $m(t)$  converges to  $\tilde{m}_b$  in  $L^2(\mu_a; K_\varepsilon)$ .

The remaining part, namely the proof that the instanton  $\tilde{m}$  is unique modulo upward translations and reflections and that it is globally stable is completely similar to the proof of the analogous properties in [14], so that we simply outline the main steps. The key ingredient is a lemma of Fife and McLeod, [19], proved for the Allen-Cahn equation. In the present context it says that given any  $m_0 \in L^\infty(K_\varepsilon; [-1, 1])$  such that

$$\liminf_{r_d \rightarrow +\infty} m_0(r) > 0, \quad \limsup_{r_d \rightarrow -\infty} m_0(r) < 0 \tag{C.35}$$

there are functions  $q_t$ ,  $a_t$  and  $b_t$ ,  $t \geq 0$ , so that for all  $t \geq 0$

$$\tilde{m}_{a_t} - q_t \leq m_t \leq \tilde{m}_{b_t} + q_t \tag{C.36}$$

where  $m_t$ ,  $t \geq 0$ , is the solution of (C.4) with initial datum  $m_0$ .

Furthermore  $q_t$ ,  $a_t$  and  $b_t$  converge exponentially fast as  $t \rightarrow +\infty$  with  $q_t \rightarrow 0$  as  $t \rightarrow +\infty$ . The proof of (C.36) is just like in [14] and very close to the original one of Fife and McLeod. With the local stability (that is already proven) and (C.36) we can now apply the same argument used in [14] to show that the only stationary solution of (4.19) in  $K_\varepsilon$  that satisfies (C.35)



are translations of the instantons. Using this and exploiting the monotonicity of the functional  $\mathcal{F}^{(1)}(\cdot; K_\varepsilon)$  one can adapt the Fife-McLeod proof of global stability of the instantons for Allen-Cahn to the present context, just as done in [14] for the  $d = 1$  case.

### Construction of the kernel $J^{(2)}(r, r')$

Let  $S_t(r, \hat{v})$ ,  $t \in \mathbb{R}$ ,  $r \in \tilde{K}_\varepsilon$ ,  $\hat{v} \in \partial B_1(0)$ , be the time flow for a point particle in  $\tilde{K}_\varepsilon$  with elastic collisions on  $\partial \tilde{K}_\varepsilon$ . Here  $(r, \hat{v})$  denotes the initial position and velocity,  $t$  the time and  $S_t(r, \hat{v}) =: (r_t, \hat{v}_t)$  the position and velocity of the particle at time  $t$ . Since  $\partial \tilde{K}_\varepsilon$  is convex there are no tangential collisions and we can conclude that, except at the collisions,  $S_t(r, \hat{v})$  is smooth and the Jacobian of the transformation  $(t, \hat{v}) \rightarrow S_t(r, \hat{v})$  is non-zero. Moreover, by the Liouville theorem, the flow  $S_t$  preserves the Lebesgue measure on  $\tilde{K}_\varepsilon \times \partial B_1(0)$ .

We call  $\lambda(d\hat{v})$  the surface measure on  $\partial B_1(0)$  and

$$p(dt d\hat{v}) := J(t) t^{d-1} dt \lambda(d\hat{v})$$

We define  $J^{(2)}(r, r')$  by setting, for any  $f \in L^\infty(\tilde{K}_\varepsilon)$  (also thought of as a function on  $\tilde{K}_\varepsilon \times \partial B_1(0)$  that does not depend on  $\hat{v}$ ) and any  $r \in \tilde{K}_\varepsilon$ ,

$$\int_{\tilde{K}_\varepsilon} dr' J^{(2)}(r, r') f(r') = \int_{[0,1] \times \partial B_1(0)} p(dt d\hat{v}) f(S_t(r, \hat{v}))$$

Property 1 of  $J^{(2)}$  follows from the fact that  $t \leq 1$  and that the speed is 1; the equality

$$\int_{[0,1] \times \partial B_1(0)} p(dt d\hat{v}) = \int_{\mathbb{R}^d} dr' J(r, r') = 1$$

proves property 3. Property 4 holds because  $S_t(r, \hat{v}) = r + t\hat{v}$  if  $d(r, \partial \tilde{K}_\varepsilon) \geq 1$ , in fact in such a case the point does not collide before  $t = 1$ . Property 5 follows from the fact that the evolution of the last coordinate  $r_d$  is like in the free motion, as the walls  $\partial \tilde{K}_\varepsilon$  are parallel to the  $r_d$ -axis.

To prove property 2 we write for  $g, f \in L^\infty(\tilde{K}_\varepsilon \times \partial B_1(0))$  which do not depend on  $\hat{v}$  but only on  $r$ :

$$\begin{aligned} & \int_{\tilde{K}_\varepsilon \times \tilde{K}_\varepsilon} dr dr' J^{(2)}(r, r') g(r) f(r') \\ &= \int_0^1 dt J(t) t^{d-1} \int_{\tilde{K}_\varepsilon} dr \int_{\partial B_1(0)} \lambda(d\hat{v}) g(r) f(S_t(r, \hat{v})) \\ &= \int_0^1 dt J(t) t^{d-1} \int_{\tilde{K}_\varepsilon} dr' \int_{\partial B_1(0)} \lambda(d\hat{v}') f(r') g(S_{-t}(r', \hat{v}')) \end{aligned}$$

having used Liouville theorem. We now observe that the position coordinate is the same in  $S_{-t}(r', \hat{v}')$  and  $S_t(r', -\hat{v}')$  and since  $g$  does not depend on the velocity we may write as well  $g(S_t(r', -\hat{v}'))$  in the last integral. By the symmetry of the measure  $d\lambda$  under the change  $\hat{v} \rightarrow -\hat{v}$  we then complete the proof of property 2. All the properties of  $J^{(2)}(r, r')$  listed in Section 4 have been proved.

## APPENDIX D

In this Appendix we recall some basic notions of geometric measure theory; we refer to the book of Evans and Gariepy, [17], for more details.

We recall that a set  $E \subset \mathcal{I}$  has finite perimeter when its characteristic function  $\mathbf{1}_E$  belongs to the space  $BV(\mathcal{I})$  of the functions on  $\mathcal{I}$  of bounded variation. When  $E$  has finite perimeter, there exists a set  $\partial^*E$  (reduced boundary of  $E$ ) and a function  $\nu : \partial^*E \rightarrow \mathbb{R}^d$ ,  $|\nu| = 1$  (generalized outer normal to  $\partial^*E$ ) such that for every vector field  $\phi \in \mathcal{C}^1(\mathcal{I})$  the following generalized form of Gauss-Green formula holds:

$$\int_E dr \operatorname{div} \phi = - \int_{\partial^*E} d\lambda \phi \cdot \nu \quad (\text{D.1})$$

where  $d\lambda$  denotes (a suitable extension of) the  $(d-1)$ -dimensional surface measure. The set  $\partial^*E$  is rectifiable in a measure theoretic sense, the precise statement may be found in Section 5.7 of [17].

The perimeter  $P(E)$  of  $E$  is given by the surface measure  $\lambda(\partial^*E)$  of its reduced boundary. When  $u \in BV(\mathcal{I}; \{\pm m_\beta\})$ , we define the perimeter  $P(u)$  of  $u$  as the perimeter of the set

$$E := \{r \in \mathcal{I} : u(r) = -m_\beta\} \quad (\text{D.2})$$

The following approximation result can be found in [8].

**Theorem D.1.** Let  $E$ ,  $\partial^*E$  and  $\nu$  be as above. Then for any  $\zeta > 0$  there exists a set  $F$  of class  $\mathcal{C}^1$  such that

$$|E \Delta F| < \zeta, \quad \lambda(\partial^*E \Delta \partial F) < \zeta \quad (\text{D.3})$$

where  $\Delta$  denotes the symmetric difference of sets.

The following result has been used in Section 4:

**Theorem D.2.** *Covering theorem.* Let  $u \in BV(\mathcal{I}; \{\pm m_\beta\})$ . Then for any  $0 < \zeta < 1$  there is  $h > 0$  and there are disjoint parallelepipeds

$R_1, \dots, R_n$  in  $\mathbb{R}^d$  with bases  $\mathbb{R}^{d-1}$ -parallelepipeds  $B_1, \dots, B_n$  respectively, and equal height  $2h$ , so that

$$\frac{1}{h} \sum_{i=1}^n \int_{R_i} dr |\chi_{R_i} - u| < \zeta; \quad \left| \sum_{i=1}^n |B_i| - P(u) \right| < \zeta \quad (\text{D.4})$$

where  $\chi_{R_i} := m_\beta(\mathbf{1}_{R_i^+} - \mathbf{1}_{R_i^-})$ .

It is possible to take all the  $R_i$  congruent to the same  $d$ -dimensional cube  $R$  of size  $2h$ .

*Proof.* (Sketch.) Using Theorem D.1 we may restrict to the case  $\partial E$  of class  $\mathcal{C}^1$ , see (D.2). In what follows we denote by  $c_i$  positive constants, possibly depending on  $d$  and  $m_\beta$  but not on  $\zeta$  and  $h$ .

We can find pairwise disjoint open subsets  $\Sigma_1, \dots, \Sigma_m$  of  $\partial E$  which cover  $\partial E$  up to a set of (surface) measure less than  $c_1\zeta$ , and so that each  $\Sigma_i$  is (congruent to) the graph of a real function  $f_i : U_i \rightarrow \mathbb{R}$  of class  $\mathcal{C}^1$ , where  $U_i \subset \mathbb{R}^{d-1}$  is a bounded open set and  $f_i$  satisfies the bound

$$|\nabla f_i| \leq c_2\zeta \quad \text{on } U_i \quad (\text{D.5})$$

See Figures 5 and 6.




Fig.5. The shaded region is a set  $E$  with finite perimeter. Shown is a covering of parts of  $\partial E$  by parallelepipeds as in Theorem D.2. Locally  $\partial E$  is the graph of a function; the segments outside  $E$  represent symbolically the domains of definition of these functions.




Fig.6. Blow up of a single piece of the covering of Fig. 5.

We choose now  $h > 0$  small enough, in such a way that the following two conditions hold:

(i) the number  $2\sqrt{d}h$  (which is the length of the diagonal of a  $d$ -dimensional cube of side  $2h$ ) is less than the distance between  $\Sigma_i$  and  $\Sigma_j$  for any  $i \neq j$ ;

(ii) for every  $i$  we cover  $U_i$  with pairwise disjoint  $(d-1)$ -dimensional cubes  $B \subset U_i$  of side  $2h$  up to a set of measure less than  $c_3\zeta/m$ .

For any  $i = 1, \dots, m$  and any  $(d-1)$ -dimensional cube  $B \subset U_i$  appearing in (ii) centered at  $x$  we construct the  $d$ -dimensional cube  $R \subset R^d$  with basis  $B$  centered at the point  $(x, f_i(x))$  and with height  $2h$ , see Figure 6. Denote by  $R_1, \dots, R_n$  (with basis  $B_1, \dots, B_n$  respectively) the collection of all these cubes. We also assume that  $R_i^-$  is in the direction of  $E$ .

Using (D.5) we then have

$$\left| \sum_{i=1}^n |B_i| - P(u) \right| \leq \sum_{i=1}^m \left( c_3 \frac{\zeta}{m} + c_4 \zeta P(\Sigma_i) \right) \leq c_5 \zeta$$

Moreover, by construction

$$\begin{aligned} \frac{1}{h} \int_{R_i} dr |\chi_{R_i} - u| &= \frac{2m\beta}{h} (|R_i^- \cap E| + |R_i^+ \cap E^c|) \\ &\leq \frac{2m\beta}{h} 2c_2 \zeta h |B_i| \leq c_6 \zeta |B_i| \end{aligned}$$

with  $E^c$  the complement of  $E$ . Finally thanks to (D.5) and (i) one can show that the  $R_i$  are pairwise disjoint, for a suitable choice of the constants  $c_i$ . ■

## ACKNOWLEDGEMENTS

We are indebted to Pierre Picco for many useful discussions and for helpful suggestions in the proof of Proposition 4.2. This research has been

partially supported by CNR (GNFM and GNSM) and by grant CEE CHRXC-CT93-0411.

## REFERENCES

- [1] G. Bellettini, M. Cassandro, and E. Presutti, *Constrained minima of non local free energy functionals*, preprint (June 1995).
- [2] J. Bricmont, J.L. Lebowitz, and C.E. Pfister, *On the surface tension of lattice systems*, Ann. N.Y. Acad. Sci. **337** (1980), 214-223.
- [3] M. Cassandro, E. Orlandi, and E. Presutti, *Interfaces and typical Gibbs configurations for one dimensional Kac potentials*, Prob. Theory Related Fields **96** (1993), 57-96.
- [4] G. Dal Maso, *An Introduction to  $\Gamma$ -Convergence*, Birkhäuser, Boston, 1993.
- [5] R. Dal Passo and P. de Mottoni, *The heat equation with a nonlocal density dependent advection term*, preprint (1991).
- [6] E. De Giorgi, F. Colombini, and L.C. Piccinini, *Frontiere Orientate di Misura Minima e Questioni Collegate*, Quaderni Sc. Norm. Sup., Pisa, 1972.
- [7] E. De Giorgi and T. Franzoni, *Su un tipo di convergenza variazionale*, Atti Accad. Naz. Lincei, Rend. Cl. Sc. Fis. Mat. Nat. (8) **58** (1975), 842-850.
- [8] T.Q. de Gromard, *Approximation forte dans  $BV(\Omega)$* , Compt. R. Acad. Sci. Paris **301** (1985), 261-264.
- [9] A. De Masi, E. Orlandi, E. Presutti, and L. Triolo, *Glauber evolution with Kac potentials: I. Mesoscopic and macroscopic limits, interface dynamics*, Nonlinearity **7** (1994), 633-696.
- [10] A. De Masi, E. Orlandi, E. Presutti, and L. Triolo, *Glauber evolution with Kac potentials: II. Fluctuations*, preprint (1994).
- [11] A. De Masi, E. Orlandi, E. Presutti, and L. Triolo, *Glauber evolution with Kac potentials: III. Spinodal decomposition*, preprint (1994).
- [12] A. De Masi, E. Orlandi, E. Presutti, and L. Triolo, *Motion by curvature by scaling non local evolution equations*, J. Stat. Phys. **73** (1993), 543-570.
- [13] A. De Masi, E. Orlandi, E. Presutti, and L. Triolo, *Stability of the interface in a model of phase separation*, Proc. R. Soc. Edinb. **124A** (1994), 1013-1022.
- [14] A. De Masi, E. Orlandi, E. Presutti, and L. Triolo, *Uniqueness of the instanton profile and global stability in non local evolution equations*, Rend. Mat. **14** (1994), 693-723.

- [15] R.L. Dobrushin, R. Kotecky, and S. Schlosman, *Wulff Construction: A Global Shape for Local Interaction*, American Mathematical Society, Providence, Rhode Island, 1992.
- [16] T. Eisele and R.S. Ellis, *Symmetry breaking and random walks for magnetic systems on a circle*, Z. Wahr. verw. Gebiete **63** (1983), 297-348.
- [17] L.C. Evans and R.F. Gariepy, *Measure Theory and Fine Properties of Functions*, CRC Press, Boca Raton, Florida, 1992.
- [18] H. Federer, *Geometric Measure Theory*, Springer-Verlag, Berlin, 1968.
- [19] P. Fife and J.B. McLeod, *The approach of solutions of nonlinear diffusion equations to travelling front solutions*, Arch. Rat. Mech. Anal. **65** (1977), 335-361.
- [20] E. Giusti, *Minimal Surfaces and Functions of Bounded Variation*, Birkhäuser, Boston, 1984.
- [21] J. Jost, *Equilibrium maps between metric spaces*, preprint (1993).
- [22] M. Kac, G. Uhlenbeck, and P.C. Hemmer, *On the van der Waals theory of vapor-liquid equilibrium: I. Discussion of a one dimensional model*, J. Math. Phys. **4** (1963), 216-228; *II. Discussion of the distribution functions*, J. Math. Phys. **4** (1963), 229-247; *III. Discussion of the critical region*, J. Math. Phys. **5** (1964), 60-74.
- [23] J. Lebowitz and O. Penrose, *Rigorous treatment of the van der Waals Maxwell theory of the liquid vapour transition*, J. Math. Phys. **7** (1966), 98-112.
- [24] S. Masini, Thesis, Fisica, Università "Roma La Sapienza" (1994).
- [25] L. Modica, *The gradient theory of phase transition and the minimal interface criterion*, Arch. Rat. Mech. and Anal. **98** (1987), 123-142.
- [26] L. Modica and S. Mortola, *Un esempio di  $\Gamma$ -convergenza*, Boll. Un. Mat. Ital. B (5) **14** (1977), 285-294.
- [27] C.E. Pfister, *Large deviations and phase separations in the two dimensional Ising model*, Helv. Physica Acta **64** (1991), 953-1054.
- [28] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, vol. I and IV*, J. Wiley, New York, 1978.
- [29] V.A. Rokhlin, *On the fundamental ideas of measure theory*, Amer. Math. Soc. Transl. (1) **10** (1962), 1-54.
- [30] L. Simon, *Lectures on Geometric Measure Theory*, Proceedings of the Centre of Mathematical Analysis, 3, Canberra, 1984.